

Infliximab Quantitation with Reflex to Antibodies to
Infliximab, Serum

Patient ID SA00106846	Patient Name SAMPLEREPORT, INFlixIMAB	Birth Date 2001-05-01	Gender F	Age 17
Order Number SA00106846	Client Order Number SA00106846	Ordering Physician CLIENT, CLIENT	Report Notes	
Account Information C7028846 DLMP Rochester		Collected 24 May 2018 12:34		

Infliximab QN with Reflex to Ab, S

Infliximab, S

Interpretation

For clinical assessment of response to therapy, infliximab should be measured at trough. When infliximab trough concentrations are greater than 5.0 mcg/mL, clinically relevant antibodies-to-infliximab are unlikely and reflex testing will not be performed.

1 SDL

Infliximab, S

SDL

7.7 mcg/mL

REFERENCE VALUE

Limit of Quantitation = 1.0 mcg/mL

Received: 24 May 2018 12:34

Reported: 24 May 2018 12:57

Laboratory Notes

- 1 This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

Performing Site Legend

Code	Laboratory	Address	Lab Director	CLIA Certificate
SDL	Mayo Clinic Laboratories - Rochester Superior Drive	3050 Superior Drive NW, Rochester MN 55901	William G. Morice M.D. Ph.D	24D1040592