

Final Report

PATIENT INFORMATION:

Name: Sample, Patient
Patient ID: 081234888
DOB: 06/05/1980
Age: 43
Gender: F

SPECIMEN INFORMATION:

Accession#: 1234 (O118000356)
Date Collected: 06/20/2023 11:35 AM
Date Accessioned: 06/21/2023 11:30 AM
Date Reported: 06/22/2023

CLINIC INFORMATION:


Organization: Rheumatology Clinic
Location Name: Rheumatology Clinic
Provider: Doctor Doctor, MD

Requisition Comments


MRN: X100461012

Current Test Result

aiSLE DX FRI Score



aiSLE DX FRI Score History



Final Report

PATIENT INFORMATION:

Name: Sample, Patient
 Patient ID: 081234888
 DOB: 06/05/1980
 Age: 43
 Gender: F

SPECIMEN INFORMATION:

Accession#: 1234 (O118000356)
 Date Collected: 06/20/2023 11:35 AM
 Date Accessioned: 06/21/2023 11:30 AM
 Date Reported: 06/22/2023

CLINIC INFORMATION:

Organization: Rheumatology Clinic
 Location Name: Rheumatology Clinic
 Provider: Doctor Doctor, MD

Test Description

The aiSLE*DX Flare Risk Index (FRI) test measures the concentrations of 11 blood plasma proteins. A proprietary weighted algorithm is used to calculate a flare risk score that ranges from -30 to +30. The magnitude of the score indicates the likelihood of the patient experiencing a lupus flare in the next 12 weeks.

Decision curve analysis was used to stratify the risk levels as: Low (< - 6.4), Medium (- 6.4 to 9.0), and High (>9.0). Applying these cut-offs:

- A maximum sensitivity of 97% and maximum specificity of 98% was observed for the FRI.
- Additional complementary analyses showed that a patient with a positive score is over 5 times more likely to flare in the next 12 weeks than a patient with a negative score (Odds Ratio = 5.2, p=0.0001).
- A patient with a positive score greater than 9.0 is highly likely to flare (Odds Ratio = 9.3, p=0.0233) while a patient with a negative score less than - 6.4 is not likely to flare (Odds Ratio = 18.0, p = 0.0003).

Test results are intended to aid in the assessment of risk of flare in lupus patients when used in conjunction with standard clinical assessment.

Munroe ME, et al. A Flare Risk Index Informed by Select Immune Mediators in Systemic Lupus Erythematosus. Arthritis Rheumatol. 2023 May;75(5):723-735.

* aiSLE is a registered trademark of Progentec Diagnostics, Inc.

Final Report

Progentec Diagnostics, Inc.
655 Research Parkway, Suite #538
Oklahoma City, OK 73104
Phone: (833) 587-8739
Fax: (833) 335-3314
CLIA#: 37D2153168 CAP#: 8352739

PATIENT INFORMATION:

Name: Sample, Patient
Patient ID: 081234888
DOB: 06/05/1980
Age: 43
Gender: F

SPECIMEN INFORMATION:

Accession#: 1234 (O118000356)
Date Collected: 06/20/2023 11:35 AM
Date Accessioned: 06/21/2023 11:30 AM
Date Reported: 06/22/2023

CLINIC INFORMATION:

Organization: Rheumatology Clinic
Location Name: Rheumatology Clinic
Provider: Doctor Doctor, MD

aiSLE DX FRI Cytokine Panel Summary

Biomarker Name	Results (pg/mL)	Ref. Interval* (pg/mL)	Flag**
01) Monocyte chemoattractant protein-1 (MCP1)/CCL2	82	54-368	Normal
02) Interleukin-5 (IL-5)	0.350	0.260-2.000	Normal
03) Interleukin-17A (IL-17A)	3.00	2.100-11.000	Normal
04) Interleukin-7 (IL-7)	1.60	1.10-12.00	Normal
05) Interleukin-4 (IL-4)	2.50	1.000-3.200	Normal
06) Tumor necrosis factor-a (TNF-a)	3.90	1.80-12.00	Normal
07) Monocyte chemoattractant protein-3 (MCP3)/CCL7	4.60	0.82-16.00	Normal
08) B Lymphocyte Stimulator (BAFF)/ BlyS	650	370-995	Normal
09) Osteopontin (OPN)	50000	14958-95972	Normal
10) Tumor necrosis factor receptor 1 (TNFR1)/TNFRSF1A	1000	617-1595	Normal
11) Tumor necrosis factor receptor 2 (TNFR2)/TNFRSF1B	2200	1079-3589	Normal
aiSLE DX FRI Score	-17.1		Normal

*Reference intervals were determined from the analysis of 120 apparently healthy control individuals consisting of European Americans (66%), African Americans (23%) and other races and ethnicities (11%) analyzed in Progentec Diagnostics laboratory. **High, Normal and Low is provided as a guide relative to this healthy control population measured with this immunoassay method.

Clinical interpretation of individual biomarker levels has not been established but may be indicative of the pathophysiology of immune, infectious, or inflammatory pathologies.

Progentec Diagnostics testing uses an immunoassay to measure the concentrations of plasma proteins. Progentec Diagnostics developed and characterized this test. It is intended for clinical use and the reported results should be interpreted in relation to each patient's clinical condition and medical history. This test is a laboratory developed test (LDT) and is not cleared by the US Food and Drug Administration (FDA). FDA clearance is not currently required for LDTs. Progentec Diagnostics Clinical Laboratory is certified under the Clinical Laboratory Improvement Amendment (CLIA) of 1988 and accredited by the College of American Pathologists to perform high complexity clinical testing.