

Patient ID SA00175021	Patient Name TESTINGRNV, ABNORMAL	Birth Date 1985-11-11	Sex F	Age 39
Order Number SA00175021	Client Order Number SA00175021	Ordering Physician CLIENT,CLIENT	Report Notes	
Account Information C7028846 DLMP Rochester		Collected 28 Mar 2025 00:00		

TP53 Mutation Analysis, Tumor

Result

MCR

Provided diagnosis: lymph node, mantle cell lymphoma, pleomorphic subtype

The following CLINICALLY RELEVANT VARIANT was detected:

Gene: TP53

DNA Change: c.402T>A (Exon 5)

Amino Acid Change: p.F134L (Phe134Leu)

Variant Allele Frequency: 77.9%

No other reportable sequence variants were detected within the analyzed regions of the tested gene(s) listed in the method description.

Interpretation

MCR

TP53 c.402T>A (p.F134L) (Exon 5)

The TP53 gene encodes the tumor suppressor p53, a protein that is involved in the DNA damage cell cycle checkpoint and causes cell cycle arrest when it senses DNA damage. p53 can also activate DNA repair genes or induce apoptosis in the presence of DNA damage. Alterations in TP53 typically result in a loss of p53 protein function. Additionally, some alterations are predicted to have oncomorphic or gain-of-function properties leading to downstream oncogenic effects, including genomic instability and excessive cell proliferation [PMID:9039259, PMID:15625370, PMID:11400116, PMID:12826609, PMID:21760960, PMID:18802452].

The presence of TP53 mutations in both solid tumors and hematologic malignancies is associated with tumor aggressiveness and poor survival [PMID: 35444210, PMID: 16861262, PMID: 2887079, PMID: 2297721, PMID: 35561299].

There are currently no known clinically approved therapies that specifically target TP53 mutations.

Additional Information

MCR

CLINICAL TRIALS

Possible clinical trials of benefit for this patient can be found at the following sites:

1) ClinicalTrials.gov:

www.clinicaltrials.gov/ct2/search/advanced

2) Mayo Clinic: www.mayo.edu/research/clinical-trials/ 3) National Cancer Institute: www.cancer.gov/clinicaltrials/search

REFERENCE TRANSCRIPT

Sequence variant nomenclature is based on the following RefSeq accession number (build GRCh37 (hg19)):TP53 NM_000546.

Specimen

MCR

Tissue, Tumor

Tissue ID

MCR

12345

Method

MCR

Microscopic examination is performed by a pathologist to identify areas of tumor for enrichment by macrodissection. DNA is extracted from FFPE or cytology slides, and next generation sequencing is performed to evaluate the presence of a mutation in all coding regions and exon/intron boundaries of the TP53 gene.

Variant nomenclature is based on build GRCh37 (hg19). For details about gene transcripts (RefSeq accession numbers), specific targeted regions of each gene, and additional information on this test, see www.mayocliniclabs.com (Test ID TP53).

Disclaimer

1 MCR

This test cannot differentiate between somatic and germline alterations. Additional testing may be necessary to clarify the significance of results if there is a potential hereditary risk.

DNA variants of uncertain significance may be identified.

Performing Site Legend

Code	Laboratory	Address	Lab Director	CLIA Certificate
MCR	Mayo Clinic Laboratories - Rochester Main Campus	200 First Street SW, Rochester, MN 55905	Nikola Baumann Ph.D.	24D0404292



Patient ID SA00175021	Patient Name TESTINGRNV, ABNORMAL	Birth Date 1985-11-11	Sex F	Age 39
Order Number SA00175021	Client Order Number SA00175021	Ordering Physician CLIENT,CLIENT	Report Notes	
Account Information C7028846 DLMP Rochester		Collected 28 Mar 2025 00:00		

A negative result does not rule out the presence of a variant that may be present but below the limits of detection of this assay. The analytical sensitivity of this assay for sequence reportable alterations is 5% mutant allele frequency with a minimum coverage of 500X in a sample with at least 20% tumor content.

Point mutations and small insertion/deletion mutations will be detected in the TP53 gene only. This test may detect single exon deletions but does not detect multi-exon deletions, duplications, larger-scale genomic copy number variants, copy neutral loss of heterozygosity (LOH), or epigenetic modifications such as promoter methylation. DELINS of 1,000 bp or less are detectable with at least 50 or more supporting reads.

Variant allele frequency (VAF) is the percentage of sequencing reads supporting a specific variant divided by the total sequencing reads at that position. In somatic testing, VAF should be interpreted in the context of several factors including, but not limited to: tumor purity/heterogeneity/copy number status (ploidy, gains/losses, loss of heterozygosity) and sequencing artifact/misalignment [PMID: 27144058, PMID: 29769535].

Rare polymorphisms may be present that could lead to

false-negative or false-positive results.

The presence or absence of a variant may not be predictive of response to therapy in all patients.

Test results should be interpreted in the context of clinical, tumor sampling, histopathological, and other laboratory data. If results obtained do not match other clinical or laboratory findings, contact the laboratory for discussion. Misinterpretation of results may occur if the information provided is inaccurate and/or incomplete.

Reliable results are dependent on adequate specimen collection and processing. This test has been validated on cytology slides and formalin-fixed, paraffin-embedded tissues; other types of fixatives are discouraged. Improper treatment of tissues, such as decalcification, may cause PCR failure.

Released By

Sounak Gupta, M.B.B.S., Ph.D.

MCR

Received: 29 Mar 2025 17:04

Reported: 06 Apr 2025 11:31

Laboratory Notes

- 1 This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

Performing Site Legend

Code	Laboratory	Address	Lab Director	CLIA Certificate
MCR	Mayo Clinic Laboratories - Rochester Main Campus	200 First Street SW, Rochester, MN 55905	Nikola Baumann Ph.D.	24D0404292