



PATIENT NAME TESTING, GNF7_ABNORMAL				ORDER NUMBER M701000586
PATIENT ID SA00157704	DATE OF BIRTH 02/03/2004	AGE 18 Y	SEX Female	REQUESTED BY CLIENT CLIENT
COLLECTED 2/1/2023, 12:00 AM	RECEIVED 2/2/2023, 2:57 PM	REPORTED 2/27/2023, 10:44 AM		
The collected, received, and reported dates and times on the report are in the time zone of the performing location. 7028846 MCL RochesterCampus Rochester MN 55901				CLIENT ORDER NUMBER SA00157704 CLIENT MRN SA00157704

TEST DESCRIPTION

Evaluation of the F7 gene associated with factor VII deficiency.

SPECIMEN

WB Whole Blood

RESULT SUMMARY

Pathogenic Variant Detected

RESULT

Gene (Transcript)	Variant	Zygosity	Classification
F7 NM_000131.4	p.Pro464Hisfs*32 c.1391del Chr13:g.113773312del	heterozygous	PATHOGENIC

The following heterozygous PATHOGENIC variant was detected:

F7 (NM_000131.4), chr13(GRCh37):g.113773312del, c.1391del, p.Pro464Hisfs*32 (p.P464Hfs*32)

INTERPRETATION

F7 c.1391del (p.Pro464Hisfs*32), PATHOGENIC

The heterozygous c.1391del (p.Pro464Hisfs*32) deletion in the F7 gene (MIM:613878) is classified as pathogenic. Pathogenic variants in the F7 gene have been associated with autosomal recessive factor VII deficiency. This variant has been reported in several unrelated individuals with a clinical diagnosis of factor VII deficiency (1-6). Furthermore, functional studies have been performed to support the pathogenicity of this variant. This variant has been shown to decrease protein secretion (2,7-9) may also slow protein synthesis (7) or increase protein instability or degradation (2). This alteration is observed in the large population genetics database gnomAD in the heterozygous state in 1 out of 3529 individuals in the subpopulation Other (minor allele frequency [MAF] = 0.0142%) and at a lower frequency (MAF = 0.0136%) in the European (non-Finnish) subpopulation (10,11). This result is supportive of carrier status for autosomal recessive factor VII deficiency. However, the presence of an undetectable second variant cannot be ruled out. Clinical correlation is recommended.

Individuals may have a pathogenic variant in the F7 gene that is not detectable by the methods utilized. Additionally, pathogenic variants in other genes not interrogated by this assay may cause a similar phenotype.

This result should be interpreted in the context of clinical findings, family history, and other laboratory testing.

Consultation with a genetics professional is recommended for interpretation of this result and to determine whether



1-800-533-1710

F7 Gene, Full Gene NGS

GNF7

PATIENT NAME TESTING, GNF7_ABNORMAL					ORDER NUMBER M701000586
PATIENT ID SA00157704	DATE OF BIRTH 02/03/2004	AGE 18 Y	SEX Female	REQUESTED BY CLIENT CLIENT	
COLLECTED 2/1/2023, 12:00 AM	RECEIVED 2/2/2023, 2:57 PM	REPORTED 2/27/2023, 10:44 AM			
The collected, received, and reported dates and times on the report are in the time zone of the performing location.				CLIENT ORDER NUMBER SA00157704	
7028846 MCL RochesterCampus Rochester MN 55901				CLIENT MRN SA00157704	

reproductive risk assessment and familial testing may be of benefit to this family. Genetic testing for family members is available by ordering FMTT / Familial Mutation, Targeted Testing for the specific variant(s) detected. Please contact the laboratory at 1-800-533-1710 or the online test catalog at www.mayocliniclabs.com for information about FMTT.

REFERENCES:

- 1) Herrmann FH, Wulff K, Auberger K, et al. Molecular biology and clinical manifestation of hereditary factor VII deficiency. *Semin Thromb Hemost.* 2000;26(4):393-400. doi:10.1055/s-2000-8458 (PMID 11092214)
- 2) Fromovich-Amit Y, Zivelin A, Rosenberg N, Tamary H, Landau M, Seligsohn U. Characterization of mutations causing factor VII deficiency in 61 unrelated Israeli patients. *J Thromb Haemost.* 2004;2(10):1774-1781. doi:10.1111/j.1538-7836.2004.00921.x (PMID 15456489)
- 3) Mariani G, Herrmann FH, Dolce A, et al. Clinical phenotypes and factor VII genotype in congenital factor VII deficiency. *Thromb Haemost.* 2005;93(3):481-487. doi:10.1160/TH04-10-0650 (PMID 15735798)
- 4) Herrmann FH, Wulff K, Auerswald G, et al. Factor VII deficiency: clinical manifestation of 717 subjects from Europe and Latin America with mutations in the factor 7 gene. *Haemophilia.* 2009;15(1):267-280. doi:10.1111/j.1365-2516.2008.01910.x (PMID 18976247)
- 5) Batorova A, Mariani G, Kavakli K, et al. Inhibitors to factor VII in congenital factor VII deficiency. *Haemophilia.* 2014;20(2):e188-e191. doi:10.1111/hae.12376 (PMID 24533960)
- 6) Arbin AA, Bodkin D, Lopaciuk S, Bauer KA. Molecular analysis of Polish patients with factor VII deficiency. *Blood.* 1994;84(7):2214-2220. (PMID 7919338)
- 7) Chollet ME, Andersen E, Skarpen E, et al. Factor VII deficiency: Unveiling the cellular and molecular mechanisms underlying three model alterations of the enzyme catalytic domain. *Biochim Biophys Acta Mol Basis Dis.* 2018;1864(3):660-667. doi:10.1016/j.bbdis.2017.12.016 (PMID 29246447)
- 8) Andersen E, Chollet ME, Myklebust CF, et al. Activation of Endoplasmic Reticulum Stress and Unfolded Protein Response in Congenital Factor VII Deficiency. *Thromb Haemost.* 2018;118(4):664-675. doi:10.1055/s-0038-1637690 (PMID 29618153)
- 9) Branchini A, Rizzotto L, Mariani G, et al. Natural and engineered carboxy-terminal variants: decreased secretion and gain-of-function result in asymptomatic coagulation factor VII deficiency. *Haematologica.* 2012;97(5):705-709. doi:10.3324/haematol.2011.049403 (PMID 22180436)
- 10) dbSNP: www.ncbi.nlm.nih.gov/snp; Sherry ST, Ward M and Sirotkin K. dbSNP-Database for Single Nucleotide Polymorphisms and Other Classes of Minor Genetic Variation. *Genome Res.* 1999;9:677-679 (PMID 10447503)
- 11) gnomAD Browser: gnomad.broadinstitute.org/; Karczewski K, Francioli LC, MacArthur DG, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature.* 2020; 581(7809):434-443 (PMID 32461654)

METHOD

Next generation sequencing (NGS) and/or Sanger sequencing was performed to test for the presence of variants in coding regions and intron/exon boundaries of the gene analyzed. The human genome reference GRCh37/hg19 build was



1-800-533-1710

F7 Gene, Full Gene NGS

GNF7

PATIENT NAME TESTING, GNF7_ABNORMAL				ORDER NUMBER M701000586
PATIENT ID SA00157704	DATE OF BIRTH 02/03/2004	AGE 18 Y	SEX Female	REQUESTED BY CLIENT CLIENT
COLLECTED 2/1/2023, 12:00 AM	RECEIVED 2/2/2023, 2:57 PM	REPORTED 2/27/2023, 10:44 AM		
The collected, received, and reported dates and times on the report are in the time zone of the performing location.				CLIENT ORDER NUMBER SA00157704
7028846 MCL RochesterCampus Rochester MN 55901				CLIENT MRN SA00157704

used for sequence read alignment. At least 99% of the bases are covered at a read depth >30X. Sensitivity is estimated at >99% for single nucleotide variants, >94% for indels up to 39 base pairs, >95% for deletions up to 75 base pairs and insertions up to 47 base pairs. NGS and/or a PCR-based quantitative method was performed to test for the presence of deletions and duplications in the gene analyzed. See the Genes Analyzed field for a list of gene(s) tested.

There may be regions of genes that cannot be effectively evaluated for sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high GC content, and repetitive sequences. Confirmation of select reportable variants was performed by alternate methodologies based on internal laboratory criteria. See www.mayocliniclabs.com (TEST ID GNF7) for details regarding genes with regions not routinely covered.

GENES ANALYZED

F7

DISCLAIMER**Clinical Correlations**

An online research opportunity called GenomeConnect (genomeconnect.org), a project of ClinGen, is available for the recipient of this genetic test. This patient registry collects de-identified genetic and health information to advance the knowledge of genetic variants. Mayo Clinic is a collaborator of ClinGen. This may not be applicable for all tests.

If testing was performed because of a clinically significant family history it is often useful to first test an affected family member. Detection of a reportable variant(s) in an affected family member would allow for more informative testing of at risk individuals.

To discuss the availability of further testing options or for assistance in the interpretation of these results, Mayo Clinic Laboratory genetic counselors can be contacted at 1-800-533-1710.

Technical Limitations

Next generation sequencing may not detect all types of genomic variants. In rare cases, false negative or false positive results may occur. The depth of coverage may be variable for some target regions, but assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated for sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high GC content, and repetitive sequences. Confirmation of select reportable variants was performed by alternate methodologies based on internal laboratory criteria.

Additionally, low level mosaic variants may not be detected.



1-800-533-1710
F7 Gene, Full Gene NGS

GNF7

PATIENT NAME TESTING, GNF7_ABNORMAL				ORDER NUMBER M701000586
PATIENT ID SA00157704	DATE OF BIRTH 02/03/2004	AGE 18 Y	SEX Female	REQUESTED BY CLIENT CLIENT
COLLECTED 2/1/2023, 12:00 AM	RECEIVED 2/2/2023, 2:57 PM	REPORTED 2/27/2023, 10:44 AM		
The collected, received, and reported dates and times on the report are in the time zone of the performing location.				CLIENT ORDER NUMBER SA00157704
7028846 MCL RochesterCampus Rochester MN 55901				CLIENT MRN SA00157704

This test is not designed to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Reclassification of Variants Policy

See www.mayocliniclabs.com (TEST ID GNF7) for information regarding the laboratory's policy for reclassification of variants.

Variant Evaluation

Variant curation is performed using published ACMG-AMP recommendations as a guideline. Other gene-specific guidelines may also be considered. Variants classified as benign or likely benign are not reported.

Results from in silico evaluation tools may change over time and should be interpreted with caution and professional clinical judgment.

TEST CLASSIFICATION

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the U.S. Food and Drug Administration.

RELEASED BY

SHANDA VOELTZ

Code : MCR Laboratory : Mayo Clinic Laboratories - Rochester Main Campus
Lab Director : WILLIAM MORICE MD, PHD CLIA Certificate : 24D0404292

Address : 200 FIRST STREET SW
ROCHESTER MN 55905