

Reporting Title: Vancomycin, Peak, S
Performing Location: Rochester

Ordering Guidance:

- In addition to this peak assay, both trough level and random testing are available.
1. Serum for a trough level should be drawn no more than 30 minutes prior to next dose; order VANTA / Vancomycin, Trough, Serum.
 2. Serum for random testing should be ordered as VANRA / Vancomycin, Random, Serum.

Specimen Requirements:

Collection Container/Tube: Red top
Submission Container/Tube: Plastic vial
Specimen Volume: 0.5 mL

Collection Instructions:

1. Serum for a peak level should be drawn 1 hour after completion of dose.
2. Centrifuge and aliquot serum into a plastic vial within 2 hours of collection.

Forms:

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	14 days	
	Frozen	365 days	
	Ambient	48 hours	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
VANPA	Vancomycin, Peak, S	Numeric	mcg/mL	4090-7

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80202

Reference Values:

Therapeutic: 20.0-45.0 mcg/mL