

Test Definition: APTTP

Activated Partial Thromboplastin Time, Plasma

Reporting Title: Activated Partial Thrombopl Time, P

Performing Location: Rochester

Ordering Guidance:

The primary method for therapeutic heparin monitoring is the heparin anti-Xa assay. Order HEPTP / Heparin Anti-Xa Assay, Plasma.

Specimen Requirements:

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL **Collection Instructions:**

1. Centrifuge, remove plasma and centrifuge plasma again.

2. Aliquot plasma into plastic vial leaving 0.25 mL in the bottom of centrifuged vial.

Additional Information: Double-centrifuged specimen is critical for accurate results as platelet contamination may cause

spurious results.

Specimen Type	Temperature	Time	Special Container	
Plasma Na Cit	Frozen (preferred)	30 days		
	Ambient	4 hours		

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
APTTP	Activated Partial Thrombopl Time, P	Numeric	sec	14979-9

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

85730

Reference Values:

25-37 seconds