

Reporting Title: VWD 2N (Normandy), P
Performing Location: Rochester

Additional Testing Requirements:
VWAG / von Willebrand Factor Antigen, Plasma; VWACT / von Willebrand Factor Activity, Plasma; and F8A / Coagulation Factor VIII Activity Assay, Plasma are recommended to supplement results of this test.

Necessary Information:
If performed at another laboratory, forward the results of the following tests with the specimen:
-von Willebrand factor antigen
-VWF activity (ristocetin cofactor, latex immunoassay etc)
-Factor VIII activity

These results aid in the interpretation of this test.

Specimen Requirements:
Specimen Type: Platelet-poor plasma
Collection Container/Tube: Light-blue top (3.2% sodium citrate)
Submission Container/Tube: Plastic vial
Specimen Volume: 1 mL

- Collection Instructions:**
- 1. For complete instructions, see [Coagulation Guidelines for Specimen Handling and Processing](#)
 - 2. Centrifuge, transfer all plasma into a vial, and centrifuge plasma again.
 - 3. Aliquot plasma into a separate tube leaving 0.25 mL in the bottom of the centrifuged vial.
 - 4. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally, less than or equal to -40 degrees C.

- Additional Information:**
- 1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
 - 2. Each coagulation assay requested should have its own vial.

Forms:
If not ordering electronically, complete, print, and send a [Coagulation Test Request](#) (T753) with the specimen.

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	56 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
607336	VWF:FVIIIIB	Numeric	%	90919-2
607337	VWF:FVIIIIB Interpretation	Alphanumeric		48595-3

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

85246

Reference Values:

68-106%

Pediatric reference ranges have not been established for this assay but likely achieve adult reference range by 18 years of age.