

Test Definition: ZNCU

Zinc/Creatinine Ratio, Urine

Reporting Title: Zinc/Creat Ratio, U

Performing Location: Rochester

Specimen Requirements:

Only orderable as part of a profile. For more information see ZNUCR / Zinc/Creatinine Ratio, Random, Urine.

Patient Preparation: High concentrations of barium are known to interfere with most metal tests. If barium-containing

contrast media has been administered, the specimen should not be collected for at least 96 hours.

Supplies: Urine Tubes, 10 mL (T068)

Collection Container/Tube: Clean, plastic urine collection container with no metal cap or glued insert

Submission Container/Tube: Plastic urine tube or clean, plastic aliquot container with no metal cap or glued insert

Specimen Volume: 3 mL **Collection Instructions:**

1. Collect a random urine specimen.

2. See Metals Analysis Specimen Collection and Transport for complete instructions.

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
615260	Zinc/Creat Ratio, U	Numeric	mcg/g Cr	13473-4

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

84630

Reference Values:

Only orderable as part of a profile. For more information see ZNUCR / Zinc/Creatinine Ratio, Random, Urine.

Not applicable