

Test Definition: PINP

Procollagen I Intact N-Terminal, Serum

Reporting Title: Procollagen I Intact N-Terminal, S **Performing Location:** Rochester

Ordering Guidance:

This test should not be requested in patients who have recently received radioisotopes, therapeutically or diagnostically, because of potential assay interference. A recommended time period before collection cannot be provided, as it depends on the isotope administered, the dose given, and the clearance rate in the individual patient. Specimens will be screened for radioactivity prior to analysis. Radioactive specimens received in the laboratory will be held and assayed after the radioactivity has sufficiently decayed. This will result in a test delay.

Specimen Requirements:

Collection Container/Tube: Preferred: Red top Acceptable: Serum gel Submission Container/Tube: Plastic vial Specimen Volume: 0.5 mL Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Specimen Type	Temperature	Time	Special Container	
Serum	Frozen (preferred)	180 days		
	Ambient	7 days		
	Refrigerated	7 days		

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
61695	Procollagen I Intact N-Terminal, S	Numeric	mcg/L	47255-5

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information: 83519

Reference Values:

Adult male: 22-87 mcg/L

Adult female premenopausal: 19-83 mcg/L

Adult female postmenopausal: 16-96 mcg/L

Reference values have not been established for patients who are younger than 18 years of age.