

Test Definition: OXYCU

Oxycodone with Metabolite Confirmation, Random, Urine

Reporting Title: Oxycodone w/metabolite Conf, U

Performing Location: Rochester

Ordering Guidance:

For situations where chain of custody is required, a Chain-of-Custody Kit (T282) is available. For chain-of-custody testing, order OXYCX / Oxycodone with Metabolite Confirmation, Chain of Custody, Random, Urine.

Additional drug panels and specific requests are available; call 800-533-1710 or 507-266-5700.

Additional Testing Requirements:

If urine creatinine is required or adulteration of the sample is suspected, the following test should also be ordered, ADULT / Adulterants Survey, Random, Urine.

Specimen Requirements:

Supplies: Sarstedt 5 mL Aliquot Tube (T914)

Collection Container/Tube: Plastic urine container **Submission Container/Tube:** Plastic, 5-mL tube

Specimen Volume: 3 mL Collection Instructions:

- 1. Collect a random urine specimen.
- 2. No preservative.

Additional Information:

- 1. No specimen substitutions.
- 2. STAT requests are not accepted for this test.
- 3. Submitting <1 mL will compromise our ability to perform all necessary testing.

Forms:

If not ordering electronically, complete, print, and send a <u>Therapeutics Test Request</u> (T831) with the specimen.

Specimen Type	Temperature	Time	Special Container	
Urine	Refrigerated (preferred)	28 days		
	Frozen	28 days		
	Ambient	14 days		

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
62616	Oxycodone-by LC-MS/MS	Alphanumeric	ng/mL	16249-5
35965	Oxymorphone-by LC-MS/MS	Alphanumeric	ng/mL	17395-5
36022	Oxycodone Interpretation	Alphanumeric		69050-3
42010	Noroxycodone-by LC-MS/MS	Alphanumeric	ng/mL	61425-5
42011	Noroxymorphone-by LC-MS/MS	Alphanumeric	ng/mL	90894-7

LOINC® and CPT codes are provided by the performing laboratory.



Test Definition: OXYCU

Oxycodone with Metabolite Confirmation, Random, Urine

Supplemental Report:

No

CPT Code Information:

80365

G0480 (if appropriate)

Reference Values:

Negative

Cutoff concentrations by liquid chromatography-tandem mass spectroscopy:

Oxycodone: 25 ng/mL Noroxycodone: 25 ng/mL Oxymorphone: 25 ng/mL Noroxymorphone: 25 ng/mL