

Reporting Title: Chromium Occupat Exp, Random, U
Performing Location: Rochester

Ordering Guidance:
High concentrations of gadolinium and iodine are known to interfere with most metal tests. If either gadolinium- or iodine-containing contrast media has been administered, a specimen should not be collected for 96 hours.

Specimen Requirements:
Supplies: Sarstedt Aliquot Tube 5 mL (T914)
Collection Container/Tube: Clean, plastic urine collection container
Submission Container/Tube: Plastic, 5-mL tube or a clean, plastic aliquot container with no metal cap or glued insert
Specimen Volume: 3 mL
Collection Instructions:
1. Collect a random urine specimen.
2. See [Metals Analysis Specimen Collection and Transport](#) for complete instructions.

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	28 days	
	Frozen	28 days	
	Ambient	14 days	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
CRETR	Creatinine, Random, U	Numeric	mg/dL	2161-8
607761	Chromium Occupational Exposure	Numeric	mcg/g Cr	13464-3
608390	Chromium Concentration	Numeric	mcg/L	5623-4

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:
No

Components:

Test Id	Reporting Name	CPT Units	CPT Code	Always Performed	Available Separately
CROM1	Chromium Occupational Exposure, U	1	82495	Yes	No
CRETR	Creatinine, Random, U	1	82570	Yes	No

CPT Code Information:
82495
82570

Reference Values:

0-17 years: Not established

> or =18 years: The American Conference of Governmental Industrial Hygienists (ACGIH) Biological Exposure Index (BEI) for daily occupational exposure to hexavalent chromium in urine is an increase of 10.0 mcg/L between pre-shift and post-shift urine collections. The ACGIH BEI for long- and short-term hexavalent chromium in urine is an end-of-shift concentration above 24.9 mcg/L at the end of the work week.