

# **Test Definition: FHLC**

Immunoglobulin M (IgM) Heavy and Light Chain (HLC) Pairs, Kappa and Lambda with Ratio

**Reporting Title:** IgM Heavy Light Chains (HLC), S **Performing Location:** LabCorp Burlington

## **Specimen Requirements:**

Patient Preparation: Patient should be fasting for eight hours to avoid lipemic sample interference.

Specimen Type: Serum

Collection Container/Tube: SST or Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.75 mL

#### **Collection Instructions:**

1. Draw blood in a plain red-top tube, serum gel tube(s) is also acceptable.

2. Separate serum immediately after coagulation (30 minutes) to prevent hemolysis. Send 0.75 mL of serum refrigerated in a plastic vial.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Ambient	7 days	
	Frozen	7 days	

## **Result Codes:**

Result ID	Reporting Name	Туре	Unit	LOINC®
Z5399	lgM Kappa, S	Alphanumeric		74866-5
Z5400	lgM Lambda, S	Alphanumeric		74867-3
Z5401	IgM K/L HLC Ratio	Alphanumeric		74870-7

LOINC<sup>®</sup> and CPT codes are provided by the performing laboratory.

#### Supplemental Report:

No

# **CPT Code Information:**

83883 x 2

Reference Values:

IgM Kappa (g/L): 0.29-1.82 IgM Lambda (g/L): 0.17-0.94 IgM Kappa:IgM Lambda ratio: 0.96-2.30