

Test Definition: 170HP

17-Hydroxypregnenolone, Serum

Reporting Title: 17-Hydroxypregnenolone, S

Performing Location: Rochester

Specimen Requirements: Collection Container/Tube:

Preferred: Red top **Acceptable:** Serum gel

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Centrifuge and aliquot serum into plastic vial.

Specimen Type	Temperature	Time	Special Container
Serum	Frozen (preferred)	28 days	
	Refrigerated	28 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
81151	17-Hydroxypregnenolone, S	Numeric	ng/dL	6765-2

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

84143

Reference Values:

CHILDREN*

Males

Premature (26-28 weeks): 1,219-9,799 ng/dL Premature (29-36 weeks): 346-8,911 ng/dL Full term (1-5 months): 229-3,104 ng/dL 6 months-364 days: 221-1,981 ng/dL

1-2 years: 35-712 ng/dL 3-6 years: <277 ng/dL 7-9 years: <188 ng/dL 10-12 years: <393 ng/dL 13-15 years: 35-465 ng/dL 16-17 years: 32-478 ng/dL

TANNER STAGES Stage I: <209 ng/dL



Test Definition: 170HP

17-Hydroxypregnenolone, Serum

Stage II: <356 ng/dL Stage III: <451 ng/dL Stage IV-V: 35-478 ng/dL

Females

Premature (26-28 weeks): 1,219-9,799 ng/dL Premature (29-36 weeks): 346-8,911 ng/dL Full term (1-5 months): 229-3,104 ng/dL 6 months-364 days: 221-1,981 ng/dL

1-2 years: 35-712 ng/dL 3-6 years: <277 ng/dL 7-9 years: <213 ng/dL 10-12 years: <399 ng/dL 13-15 years: <408 ng/dL 16-17 years: <424 ng/dL

TANNER STAGES

Stage I: <236 ng/dL Stage II: <368 ng/dL Stage III: <431 ng/dL Stage IV-V: <413 ng/dL

ADULTS

Males

> or =18 years: 55-455 ng/dL

Females

> or =18 years: 31-455 ng/dL

*Kushnir MM, Rockwood AL, Roberts WL, et al: Development and performance evaluation of a tandem mass spectrometry assay for 4 adrenal steroids. Clin Chem. 2006 Aug;52(8):1559-1567

To convert to nmol/L, multiply the value in ng/dL by 0.03159757.