

Test Definition: IGA

Immunoglobulin A (IgA), Serum

Reporting Title: Immunoglobulin A (IgA), S

Performing Location: Rochester

Ordering Guidance:

Cascade testing is recommended for celiac disease. Cascade testing ensures that testing proceeds in an algorithmic fashion. The following cascades are available; select the appropriate one for your specific patient situation.

- -CDCOM / Celiac Disease Comprehensive Cascade, Serum and Whole Blood: Complete testing including HLA DQ
- -CDSP / Celiac Disease Serology Cascade, Serum: Complete serology testing excluding HLA DQ
- -CDGF / Celiac Disease Gluten-Free Cascade, Serum and Whole Blood: For patients already adhering to a gluten-free diet To order individual tests, see <u>Celiac Disease Diagnostic Testing Algorithm</u>.

Specimen Requirements:

Collection Container/Tube:

Preferred: Serum gel **Acceptable:** Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms:

If not ordering electronically, complete, print, and send a <u>Gastroenterology and Hepatology Test Request</u> (T728) with the specimen.

Specimen Type	Temperature	Time	Special Container	
Serum	Refrigerated (preferred)	28 days		
	Frozen	28 days		
	Ambient	14 days		

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
IGA	Immunoglobulin A (IgA), S	Numeric	mg/dL	2458-8

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

82784

Reference Values:

0-<5 months: 7-37 mg/dL 5-<9 months: 16-50 mg/dL



Test Definition: IGA

Immunoglobulin A (IgA), Serum

9-<15 months: 27-66 mg/dL 15-<24 months: 36-79 mg/dL 2-<4 years: 27-246 mg/dL 4-<7 years: 29-256 mg/dL 7-<10 years: 34-274 mg/dL 10-<13 years: 42-295 mg/dL 13-<16 years: 52-319 mg/dL 16-<18 years: 60-337 mg/dL > or =18 years: 61-356 mg/dL