

Test Definition: INHAB

Inhibin A and B, Tumor Marker, Serum

Reporting Title: Inhibin A and B, Tumor Marker, S

Performing Location: Rochester

Specimen Requirements:

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Collection Container/Tube:

Preferred: Serum gel **Acceptable:** Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1.1 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms:

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	90 days	
	Ambient	7 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
INHA	Inhibin A, Tumor Marker, S	Numeric	pg/mL	23883-2
88722	Inhibin B, S	Numeric	pg/mL	56940-0

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

Components:

Test Id	Reporting Name	CPT Units	CPT Code	Always Performed	Available Separately
INHA	Inhibin A, Tumor Marker, S	1	86336	Yes	Yes
INHB	Inhibin B, S	1	83520	Yes	Yes

CPT Code Information:

83520-Inhibin B 86336-Inhibin A

Reference Values:

INHIBIN A, TUMOR MARKER



Test Definition: INHAB

Inhibin A and B, Tumor Marker, Serum

Males: <5.0 pg/mL

Females

<11 years: <5.0 pg/mL 11-17 years: <98 pg/mL Premenopausal: <98 pg/mL Postmenopausal: <5.0 pg/mL

INHIBIN B

Males

<15 days: 68-373 pg/mL 15-180 days: 42-516 pg/mL 6 months-7 years: 24-300 pg/mL

8-30 years: 47-383 pg/mL 31-72 years: <358 pg/mL >72 years: Not established

Females

< or =12 years: <183 pg/mL</pre>

13-41 years regular Cycle (Follicular Phase): <224 pg/mL 42-51 years regular Cycle (Follicular Phase): <108 pg/mL 13-51 years regular Cycle (Luteal Phase): <80 pg/mL

>51 years (Postmenopausal): <12 pg/mL