

Test Definition: DIG

Digoxin, Serum

Reporting Title: Digoxin, S **Performing Location:** Rochester

Specimen Requirements:

Patient Preparation: For 12 hours before specimen collection do not take multivitamins or dietary supplements containing biotin (vitamin B7), which is commonly found in hair, skin, and nail supplements and multivitamins. Collection Container/Tube:

- Preferred: Serum gel
- Acceptable: Red top
- Submission Container/Tube: Plastic vial
- Specimen Volume: 1 mL
- Collection Instructions:
- 1. Draw blood 6 to 8 hours after the last dose of digoxin.
- 2. Serum gel tubes should be centrifuged within 2 hours of collection.
- 3. Red-top tubes should be centrifuged, and the serum aliquoted into a plastic vial within 2 hours of collection.

Forms:

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen: -<u>Cardiovascular Test Request Form</u> (T724)

-<u>Therapeutics Test Request</u> (T831)

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	7 days	
	Frozen	180 days	

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
DIG	Digoxin, S	Numeric	ng/mL	83093-5

LOINC[®] and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information: 80162

Reference Values:

<16 years:

Therapeutic ranges have not been established for patients who are less than 16 years of age.

> or =16 years:

Therapeutic range: 0.6-1.2 ng/mL



Test Definition: DIG

Digoxin, Serum

Toxic concentration: > or =4.0 ng/mL