

Reporting Title: Procainamide and NAPA, S
Performing Location: Rochester

Specimen Requirements:

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions:

- 1. Serum gel tubes should be centrifuged within 2 hours of collection.
- 2. Red-top tubes should be centrifuged and the serum aliquoted into a plastic vial within 2 hours of collection.

Forms:

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Type	Temperature	Time	Special Container
Serum	Frozen (preferred)	14 days	
	Refrigerated	7 days	
	Ambient	24 hours	

Result Codes:

Result ID	Reporting Name	Type	Unit	LOINC®
8683	Procainamide, S	Numeric	mcg/mL	3982-6
2461	N-acetylprocainamide, S	Numeric	mcg/mL	3834-9
2462	Procainamide + NAPA	Alphanumeric	mcg/mL	3983-4

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

80192

Reference Values:

Procainamide

Therapeutic: 4.0-10.0 mcg/mL

Critical value: >12.0 mcg/mL

N-acetylprocainamide

Therapeutic: 12.0-18.0 mcg/mL
Critical value: > or =40.0 mcg/mL