

Test Definition: VMA

Vanillylmandelic Acid, 24 Hour, Urine

Reporting Title: Vanillylmandelic Acid, 24 Hr, U

Performing Location: Rochester

Ordering Guidance:

In the past, this test has been used to screen for pheochromocytoma. However, vanillylmandelic acid (VMA) is not the analyte of choice to rule out a diagnosis of pheochromocytoma. Recommended tests for that purpose include:

- -PMET / Metanephrines, Fractionated, Free, Plasma
- -METAF / Metanephrines, Fractionated, 24 Hour, Urine
- -CATU / Catecholamine Fractionation, Free, 24 Hour, Urine

Necessary Information:

- 1. Patients age is required.
- 2. Collection duration and urine volume (in milliliters) are required.
- 3. All patients receiving L-dopa should be identified to the laboratory when this test is ordered.
- 4. Bactrim may interfere with detection of the analyte. All patients taking Bactrim should be identified to the laboratory when this test is ordered.

Specimen Requirements:

Patient Preparation: Administration of L-dopa may falsely-increase vanillylmandelic acid results; it should be discontinued 24 hours prior to and during specimen collection.

Supplies: Urine Tubes, 10 mL (T068)

Specimen Volume: 5 mL **Collection Instructions:**

- 1. Collect a 24-hour urine specimen.
- 2. Add 25 mL of 50% acetic acid as preservative at the start of collection. If specimen is refrigerated during collection, preservative may be added up to 12 hours after collection. Use 15 mL of 50% acetic acid for children <5 years old. This preservative is intended to achieve a pH of between approximately 1 and 5. If necessary, adjust urine pH to a level between 1 and 5 by adding 50% acetic acid or hydrochloric acid dropwise and checking the pH..

Additional Information: For multiple collections see <u>Urine Preservatives-Collection and Transportation for 24-Hour</u> <u>Urine Specimens.</u>

Forms:

If not ordering electronically, complete, print, and send an Oncology Test Request (T729) with the specimen.

Urine Preservative Collection Options:

Note: For addition of preservative or application of temperature information, see Specimen Required.

Ambient	No
Refrigerate	No
Frozen	No
50% Acetic Acid	Preferred
Boric Acid	ОК
Diazolidinyl Urea	No
6M Hydrochloric	ОК
Acid	
6M Nitric Acid	ОК
Sodium Carbonate	No



Test Definition: VMA

Vanillylmandelic Acid, 24 Hour, Urine

Thymol	No	
Toluene	No	
*If boric acid is used, note on specimen		
container. Also, verify that pH is in desired		
range (pH=1-5). If pH is outside of desired		
range, adjust pH with a stronger acid (acetic		
acid is preferred but other acids listed above		

could be used if available) in a dropwise fashion to bring pH into desired range.

Specimen Type	pe Temperature		Special Container
Urine	Refrigerated (preferred)		
	Frozen	180 days	

Ask at Order Entry (AOE) Questions:

	Test ID	Question ID	Description	Туре	Reportable
	VMA	TM41	Collection Duration	Plain Text	Yes
ĺ	VMA	VL39	Urine Volume	Plain Text	Yes

Result Codes:

Result ID	Reporting Name	Туре	Unit	LOINC®
3580	Vanillylmandelic Acid, Adult (>14y)	Alphanumeric	mg/24 h	3122-9
3581	Vanillylmandelic Acid, Child (<15y)	Alphanumeric	mg/g Cr	30571-4
TM41	Collection Duration	Alphanumeric		13362-9
VL39	Urine Volume	Alphanumeric		3167-4

LOINC® and CPT codes are provided by the performing laboratory.

Supplemental Report:

No

CPT Code Information:

84585

Reference Values:

<1 year: <25.0 mg/g creatinine 1 year: <22.5 mg/g creatinine 2-4 years: <16.0 mg/g creatinine 5-9 years: <12.0 mg/g creatinine 10-14 years: <8.0 mg/g creatinine

> or =15 years (adults): <8.0 mg/24 hours