

Overview

Useful For

Monitoring serum concentration of desipramine during therapy

Evaluating potential desipramine toxicity

The test may also be useful to evaluate patient compliance

Method Name

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Serum Red

Specimen Required

Collection Container/Tube: Red top (Serum gel/SST are **not acceptable**)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. Collect specimen immediately before next scheduled dose (minimum 12 hours after last dose).
2. Centrifuge and aliquot serum into plastic vial. **Serum must be separated from cells within 2 hours of collection.**

Forms

If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	28 days	
	Frozen	28 days	
	Ambient	7 days	

Clinical & Interpretive

Clinical Information

Desipramine is a tricyclic antidepressant; it also is a metabolite of imipramine. These drugs have also been employed in the treatment of enuresis (involuntary urination) in childhood and severe obsessive-compulsive neurosis. Desipramine is the antidepressant of choice in patients where maximal stimulation is indicated.

The therapeutic concentration of desipramine is 100 to 300 ng/mL. About 1 to 3 weeks of treatment are required before therapeutic effectiveness becomes apparent.

The most frequent side effects are those attributable to anticholinergic effects: dry mouth, constipation, dizziness, tachycardia, palpitations, blurred vision, and urinary retention. These occur at blood concentrations in excess of 400 ng/mL, although they may occur at therapeutic concentrations in the early stage of therapy. Cardiac toxicity (first-degree heart block) is usually associated with blood concentrations in excess of 400 ng/mL.

Reference Values

Therapeutic concentration: 100-300 ng/mL

Note: Therapeutic ranges are for specimens collected at trough (ie, immediately before next scheduled dose). Levels may be elevated in non-trough specimens.

Interpretation

Most individuals display optimal response to desipramine with serum levels of 100 to 300 ng/mL. Some individuals may respond well outside of this range or may display toxicity within the therapeutic range; thus, interpretation should include clinical evaluation. Risk of toxicity is increased with levels above 400 ng/mL.

Cautions

This test cannot be performed on whole blood. Serum must be separated from cells within 2 hours of collection; if serum is not removed within this time, tricyclic antidepressant levels may be falsely elevated due to drug release from red blood cells.

Specimens that are obtained from gel tubes are not acceptable, as the drug can absorb on the gel and lead to falsely decreased concentrations.

Clinical Reference

- Wille SM, Cooreman SG, Neels HM, Lambert WE: Relevant issues in the monitoring and toxicology of antidepressants. Crit Rev Clin Lab Sci. 2008;45(1):25-89
- Thanacoody HK, Thomas SH: Antidepressant poisoning. Clin Med. 2003;3(2):114-118

3. Hiemke C, Baumann P, Bergemann N, et al: AGNP Consensus Guidelines for Therapeutic Drug Monitoring in Psychiatry: Update 2011. Pharmacopsychiatry. 2011;44(6):195-235
4. Burtis CA, Ashwood ER, Bruns ED, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 5th ed. Elsevier; 2012

Performance

Method Description

The tricyclic antidepressants are extracted from serum using a solvent crash to precipitate proteins. The supernatant is removed and analysis is by liquid chromatography-tandem mass spectrometry (LC-MS/MS). (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Tuesday, Thursday, Sunday

Report Available

3 to 5 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80335

G0480 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
DESPR	Desipramine, S	3531-1

Result ID	Test Result Name	Result LOINC® Value
37123	Desipramine, S	3531-1