

Overview

Useful For

Evaluation of renal tubular damage

Monitoring exposure to cadmium and mercury

Method Name

AutomatedChemiluminescentImmunoMetricAssay

NY State Available

Yes

Specimen

Specimen Type

Urine

Specimen Required

Supplies: Urine Tubes, 5 mL

Container/Tube: Plastic, urine tube

Specimen Volume: 3 mL

Collection Instructions:

1. Patient should empty bladder.
2. Have patient drink at least 0.5 liters of water.
3. Within 1 hour, collect a random urine specimen.
4. Add 1 M NaOH as preservative to the collection. This preservative is intended to achieve a pH of between approximately 6 and 8.

Forms

If not ordering electronically, complete, print, and send a [Renal Diagnostics Test Request](#) (T830) with the specimen.

Specimen Minimum Volume

1 mL

Reject Due To

Specimen with pH <6	Reject
---------------------	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Frozen (preferred)	14 days	
	Refrigerated	48 hours	

Clinical and Interpretive

Clinical Information

Beta-2 microglobulin is a low-molecular-weight protein that forms the light chain component of class I histocompatibility (HLA: human leukocyte antigen) antigens.

Increased urine levels are seen in proximal tubular renal damage due to a variety of causes, including cadmium, mercury, lithium, or aminoglycoside toxicity; pyelonephritis; and Balkan nephropathy, a chronic interstitial nephritis of unknown etiology.

Reference Values

< or =300 mcg/L

Interpretation

Increased excretion is consistent with renal tubular damage.

Beta-2 microglobulin excretion is increased 100 to 1,000 times normal levels in cadmium-exposed workers.

Cautions

Degradation of beta-2 microglobulin occurs at pH <6.0.

Clinical Reference

- Ikeda M, Ezaki T, Tsukahara T, et al: Threshold levels of urinary cadmium in relation to increases in urinary beta2-microglobulin among general Japanese populations. *Toxicol Lett* 2003 Feb 3;137(3):135-141
- Moriguchi J, Ezaki T, Tsukahara T, et al: Comparative evaluation of four urinary tubular dysfunction markers, with special references to the effects of aging and correction for creatinine concentration. *Toxicol Lett* 2003 Aug 28;143(3):279-290
- Stefanovic V, Cukuranovic R, Mitic-Zlatkovic M, Hall PW: Increased urinary albumin excretion in children from families with Balkan nephropathy. *Pediatr Nephrol* 2002 Nov;17(11):913-916

Performance

Method Description

Testing is performed on the Immulite 2000. Immulite 2000 Beta-2 Microglobulin is a solid phase, 2-site chemiluminescent enzyme-labeled immunometric assay. The solid-phase bead is coated with an affinity-purified murine monoclonal anti-beta-2 antibody. The serum sample and alkaline phosphatase conjugated affinity-purified goat polyclonal anti-beta-2 antibody are incubated to bind beta-2 microglobulin into an antibody sandwich complex.

The chemiluminescent substrate, a phosphate ester of adamantyl dioxetane, in the presence of alkaline phosphatase produces light proportional to the concentration of the beta-2 microglobulin in the sample.(Product Information: IMMULITE 2000 Beta-2 Microglobulin, Siemens Healthcare Diagnostics, 2012-11-05)

PDF Report

No

Day(s) and Time(s) Test Performed

Monday through Friday; 5 a.m.-3 p.m., Saturday; 6 a.m.-3 p.m.

Analytic Time

1 day

Maximum Laboratory Time

3 days

Specimen Retention Time

2 Weeks

Performing Laboratory Location

Rochester

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved or is exempt by the U.S. Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

82232

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
B2MU	Beta-2 Microglobulin, U	1953-9

Result ID	Test Result Name	Result LOINC Value
B2MU	Beta-2 Microglobulin, U	1953-9