

Overview

Useful For

Detection of the more common potential causes of abnormal bleeding (eg, factor deficiencies/hemophilia, von Willebrand disease, factor-specific inhibitors) and a simple screen to evaluate for an inhibitor or severe deficiency of factor XIII (rare)

This test is **not useful** for assessing platelet function (eg, congenital or acquired disorders such as Glanzmann thrombasthenia, Bernard-Soulier syndrome, storage pool disease, myeloproliferative disease, associated platelet dysfunction), which requires fresh platelets

Profile Information

Test ID	Reporting Name	Available Separately	Always Performed
ALBLI	Limited Bleed Prof Interp	No	Yes
PTSC	Prothrombin Time (PT), P	Yes, (order PTTT)	Yes
APTSC	Activated Partial Thrombopl Time, P	Yes, (order APTTP)	Yes
TTSC	Thrombin Time (Bovine), P	Yes	Yes
CLFIB	Fibrinogen, Clauss, P	Yes, (order FIBTP)	Yes
DIMER	D-Dimer, P	Yes, (order DDITT)	Yes
F8A	Coag Factor VIII Activity Assay, P	Yes	Yes
F_9	Coag Factor IX Assay, P	Yes	Yes
FXIII	Factor XIII(13),Scrn	No	Yes
VWAG	von Willebrand Factor Ag, P	Yes	Yes
VWACT	von Willebrand Factor Activity, P	Yes	Yes

Reflex Tests

Test ID	Reporting Name	Available Separately	Always Performed
GBETH	General Factor Bethesda Units, P	No	No
5BETH	FV Bethesda Units, P	No	No
8BETH	FVIII Bethesda Units, P	No	No
9BETH	FIX Bethesda Units, P	No	No
F8IS	Coag Factor VIII Assay Inhib Scrn,P	No	No
RIST	Ristocetin Cofactor, P	No	No
FACTV	Coag Factor V Assay, P	Yes	No
F_7	Coag Factor VII Assay, P	Yes	No



Test ID	Reporting Name	Available Separately	Always Performed
F_10	Coag Factor X Assay, P	Yes	No
F_11	Coag Factor XI Assay, P	Yes	No
F_12	Coag Factor XII Assay, P	Yes	No
F_2	Coag Factor II Assay, P	Yes	No
F9_IS	Factor IX Inhib Scrn	No	No
F5_IS	Factor V Inhib Scrn	No	No
F2_IS	Factor II Inhib Scrn	No	No
F7_IS	Factor VII Inhib Scrn	No	No
10_IS	Factor X Inhib Scrn	No	No
11_IS	Factor XI Inhib Scrn	No	No
PNP	Platelet Neutralization Procedure	No	No
PSGN	Plasminogen Activity, P	Yes	No
PAI1	PAI-1 Ag, P	Yes	No
CH8	Chromogenic FVIII, P	Yes	No
CH9	Chromogenic FIX, P	Yes	No
ATTI	Antithrombin Antigen, P	Yes	No
ATTF	Antithrombin Activity, P	Yes	No
PTMSC	PT Mix 1:1	No	No
APMSC	APTT Mix 1:1	No	No
DRV2	DRVVT Mix	No	No
DRV3	DRVVT Confirmation	No	No
STAACL	StacLOT LA, P	No	No
VWFMP	von Willebrand Factor Multimer, P	Yes, (order VWFMS)	No
DRV1	Dilute Russells Viper Venom Time, P	Yes, (order DRV11)	No
PTFIB	PT-Fibrinogen, P	No	No
SOLFM	Soluble Fibrin Monomer	No	No
RTSC	Reptilase Time, P	Yes	No
A2PI	Alpha-2 Plasmin Inhibitor, P	Yes	No

Testing Algorithm

Initial testing includes: prothrombin time (PT), activated partial thromboplastin time (APTT), thrombin time (bovine), fibrinogen, D-dimer, coagulation factor VIII activity assay, coagulation factor IX assay, von Willebrand factor (VWF) antigen, VWF activity, factor XIII screen, and limited bleed profile interpretation.

Â

If PT is >13.9 seconds, then PT mix will be performed at an additional charge.

If APTT is > or =38 seconds, then APTT mix and dilute Russell viper venom time (DRVVT) will be performed at an additional charge.

If DRVVT ratio is > or =1.20, then DRVVT mix and DRVVT confirm will be performed at an additional charge.

If thrombin time is > or =25.0 seconds, then reptilase time will be performed at an additional charge.

If fibrinogen is <150 mg/dL, or clinically indicated, then PT-fibrinogen will be performed at an additional charge.

If D-dimer is >500 ng/mL FEU, then soluble fibrin monomer will be performed at an additional charge.

If APTT mix is > or =38 seconds and thrombin time is <35.0 seconds (no evidence of heparin), then platelet neutralization procedure will be performed at an additional charge.

If VWF activity assay is <55% or VWF activity:VWF antigen ratio is abnormally increased, then VWF ristocetin cofactor activity assay will be performed at an additional charge.

If VWF antigen is <55%, the VWF activity is <55%, or the VWF activity:VWF antigen ratio is abnormal, then VWF multimer analysis will be performed at an additional charge.

If appropriate, coagulation factor assays or Staclot lupus anticoagulant will be performed at an additional charge to clarify significant abnormalities in the screen test results.

If factor result is below the normal range, the appropriate factor inhibitor screen may be performed along with the Bethesda titrating assay, at an additional charge, if inhibitor screen is positive.

See [Hemophilia Testing Algorithm](#) in Special Instructions.

Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)
- [Coagulation Patient Information](#)
- [Hemophilia Testing Algorithm](#)
- [Coagulation Profile Comparison](#)

Method Name

PTSC, APTSC, TTSC, FXIII, F8A, F_9 : Optical Clot-Based

VWAG, VWACT, DIMER: Latex Immunoassay (LIA)

CLFIB: Clauss

NY State Available

Yes

Specimen

Specimen Type

Plasma Na Cit

Advisory Information

Multiple coagulation profile tests are available. See [Coagulation Profile Comparison](#) in Special Instructions for testing that is performed with each profile.

Shipping Instructions

Send the 6 aliquots in the same shipping container.

Necessary Information

1. If priority specimen, mark request form, give reason, and request a call-back.
2. Note if patient is currently receiving anticoagulant (eg, heparin, Coumadin [warfarin]) treatment or any medication that could affect coagulation parameters.
3. Note if patient has been recently transfused.

Specimen Required**Patient Preparation:**

1. Patient should not be receiving anticoagulant treatment (eg, warfarin, heparin). Treatment with heparin causes false-positive results of in vitro coagulation testing for lupus anticoagulant. Coumadin (warfarin) treatment may impair ability to detect the more subtle varieties of lupus-like anticoagulants.
2. Patient should also not be receiving fibrinolytic agents (streptokinase, urokinase, tissue plasminogen activator: tPA).
3. If patient has been recently transfused, it is best to perform this study pretransfusion, if possible.

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: 6 Plastic vials

Specimen Volume: 6 mL in 6 plastic vials, each containing 1 mL

Collection Instructions:

1. Specimen must be collected prior to factor replacement therapy.
2. For complete instructions, see [Coagulation Guidelines for Specimen Handling and Processing](#) in Special Instructions.
3. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
4. Aliquot plasma (1-2 mL per aliquot) into 6 separate plastic vials, leaving 0.25 mL in the bottom of centrifuged vial.
5. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally, < or = -40 degrees C.

Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

Forms

1. [Coagulation Patient Information](#) (T675) in Special Instructions.
2. If not ordering electronically, complete, print, and send a [Coagulation Test Request](#) (T753) with the specimen.

Specimen Minimum Volume

4 mL in 4 plastic vials, 1 mL each

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Clinical and Interpretive

Clinical Information

Bleeding problems may be associated with a wide variety of coagulation abnormalities or may be due to problems not associated with coagulation (trauma and surgery as obvious examples). A partial listing of causes follows.

-Deficiency or functional abnormality (congenital or acquired) of any of the following coagulation proteins: fibrinogen (factor I), factor II (prothrombin), factor V, factor VII, factor VIII (hemophilia A), factor IX (hemophilia B), factor X, factor XI (hemophilia C; bleeding severity not always proportionate to factor level), factor XIII (fibrin-stabilizing factor), von Willebrand factor (VWF antigen and activity), and alpha-2 plasmin inhibitor and plasminogen activator inhibitor (PAI-I; severe deficiency in rare cases). Neither alpha-2 plasmin inhibitor nor PAI-I are included as a routine bleeding diathesis assay component, but either can be performed if indicated or requested.

-Deficiency (thrombocytopenia) or functional abnormality of platelets such as congenital (Glanzmann thrombasthenia, Bernard-Soulier syndrome, storage pool disorders, etc) and acquired (myeloproliferative disorders, uremia, drugs, etc) disorders. Platelet function abnormalities cannot be studied on mailed-in specimens.

-Specific factor inhibitors (most commonly directed against factor VIII); factor inhibitors occur in 10% to 15% of the hemophilia population and are more commonly associated with severe deficiencies of factor VIII or IX (antigen <1%). The inhibitor appears in response to transfusion therapy with factor concentrates with no correlation of occurrence and amount of therapy. Factor VIII inhibitors may occur spontaneously in the postpartum patient, with certain malignancies, in association with autoimmune disorders (eg, rheumatoid arthritis, systemic lupus erythematosus), in the elderly, and for no apparent reason.

-Other acquired causes of increased bleeding include paraproteinemia; other factor-specific inhibitors, including

those against factor V, factor XI; or virtually any of the coagulation proteins.

-Acute disseminated intravascular coagulation/intravascular coagulation and fibrinolysis (DIC/ICF), which is a fairly common cause of bleeding. Bleeding can also occur in patients with chronic ICF.

Reference Values

An interpretive report will be provided.

Interpretation

An interpretive report will be provided.

Cautions

No significant cautionary statements

Clinical Reference

Boender J, Kruip MJ, Leebeek FW: A Diagnostic Approach to Mild Bleeding Disorders. J Thromb Haemost 2016;Aug;14(8):1507-1516 doi: 10.1111/jth.13368.

Performance

Method Description

Prothrombin time, optical clot-based:

The prothrombin time (PT) assay is performed on the Instrumentation Laboratory ACL TOP. Patient sample is incubated and combined with a PT reagent containing recombinant human tissue factor, synthetic phospholipids, calcium chloride, polybrene, and buffer. The tissue thromboplastin-factor VII/VIIa complex activates factor X. Activated factor X (factor Xa) forms a complex with factor Va, calcium, and phospholipid to activate factor II (prothrombin) to thrombin. Thrombin then acts on fibrinogen (factor I) to form fibrin which clots, the time to clot formation is measured optically using a wavelength of 671 nm providing the assay endpoint (the "prothrombin time").(Package insert: HemosIL RecombiPlasTin 2G Instrumentation Laboratory Company, R0, 03/2019)

Activated partial thromboplastin time, optical clot-based:

The activated partial thromboplastin time (APTT) assay is performed on the Instrumentation Laboratory ACL TOP. Patient sample is combined and incubated with an APTT reagent containing phospholipid, a negatively charged contact factor activator, and buffer. After a specified incubation time, calcium is added to trigger the coagulation process in the mixture. Subsequently, the time to clot formation is measured optically using a wavelength of 671 nm. Mixing studies (see APMSC / Activated Partial Thromboplastin Time (APTT) Mix 1:1, Plasma) using normal pooled plasma are performed in the Special Coagulation Laboratory on samples with a prolonged APTT, to assist in discriminating between factor deficiency states and coagulation inhibitors, unless further testing is not indicated.(Package insert: HemosIL SynthASil. Instrumentation Laboratory Company, R11, 06/2017)

Thrombin time, optical clot-based:

The thrombin time (TT) assay is performed on the Instrumentation Laboratory ACL TOP. Patient sample is combined with a bovine thrombin reagent containing bovine albumin, calcium chloride, and buffer immediately triggering the coagulation process in the mixture. Time to clot formation is measured optically using a wavelength of 671 nm.(Package insert: HemosIL Thrombin Time, Instrumentation Laboratory Company, R1 12/2018)

Fibrinogen, Clauss assay:

The Clauss fibrinogen assay is performed using the HemosIL Fibrinogen-C kit on the Instrumentation Laboratory ACL TOP. Patient sample, containing fibrinogen, is mixed with reagent containing excess thrombin. The excess thrombin converts the fibrinogen in the patient sample to fibrin. The amount of time it takes to form a clot is inversely proportional to the amount of fibrinogen present in the patient sample. (Package insert: HemosIL Fibrinogen-C. Instrumentation Laboratory Company, R7, 06/2017)

D-Dimer, latex immunoassay:

The D-dimer assay is performed using the HemosIL D-Dimer HS 500 kit on the Instrumentation Laboratory ACL TOP instrument. D-dimer is assayed in plasma by adding polystyrene latex particles coated with monoclonal antibodies specific for D-dimer domain. The latex particles agglutinate in the presence of soluble fibrin degradation products (FDP) containing the D-dimer domain. The degree of agglutination is directly proportional to the concentration of D-dimer in the sample and is determined by measuring the decrease of transmitted light caused by the aggregates (turbidimetric immunoassay). (Package insert: HemosIL D-Dimer HS 500. Instrumentation Laboratory Company, 04/2018)

Factor VIII activity, optical clot-based:

The factor VIII assay is performed on the Instrumentation Laboratory ACL TOP using the activated partial thromboplastin time (APTT) method and a factor-deficient substrate. Patient sample is combined and incubated with a factor VIII-deficient substrate (normal plasma depleted of factor VIII by immunoabsorption) and an APTT reagent. After a specified incubation time, calcium is added to trigger the coagulation process in the mixture. Then the time to clot formation is measured optically at a wavelength of 671 nm. (Owen CA Jr, Bowie EJW, Thompson JH Jr: Diagnosis of Bleeding Disorders. Second edition. Little, Brown and Company, 1975; Meijer P, Verbruggen HW, Spannagi M: Chapter 33: Clotting factors and inhibitors: Assays and Interpretation. In Laboratory Hematology Practice. Edited by K Kottke-Marchant. Wiley Blackwell Publishing, 2012, pp 435-446)

Factor IX activity, optical clot-based:

The factor IX assay is performed on the Instrumentation Laboratory ACL TOP using the APTT method and a factor-deficient substrate. Patient sample is combined and incubated with a factor IX-deficient substrate (normal plasma depleted of factor IX by immunoabsorption) and an APTT reagent. After a specified incubation time, calcium is added to trigger the coagulation process in the mixture. Then the time to clot formation is measured optically at a wavelength of 671 nm. (Owen CA Jr, Bowie EJW, Thompson JH Jr: Diagnosis of Bleeding Disorders. Second edition. Little, Brown and Company, 1975; Meijer P, Verbruggen HW, Spannagi M: Chapter 33: Clotting factors and inhibitors: Assays and Interpretation. In Laboratory Hematology Practice. Edited by K Kottke-Marchant. Wiley Blackwell Publishing, 2012, pp 435-446)

Factor XIII, clot-based:

The covalent stabilization of fibrin by thrombin-activated factor XIII (XIIIa) is the final event in the coagulation of blood. Plasma factor XIII (fibrin-stabilizing factor; FSF) zymogen consists of 2 "A" and 2 "B" subunits, the "A" subunits containing an active-center sulfhydryl grouping mediating the transamidase activity of the enzyme. The action of thrombin converts fibrinogen to fibrin monomer causing the monomeric molecules to polymerize and be held together by noncovalent hydrogen bonds. These bonds can be broken by 5 M urea or weak acid solutions in the absence of factor XIII. Subsequent to fibrin polymerization by hydrogen bonding, the action of factor XIII results in the formation of covalent bonds that cannot be broken by 5 M urea or weak acid solutions as used in this procedure (1% monochloroacetic acid). Dissolution of a clot by urea or monochloroacetic acid is therefore a qualitative test for factor XIII activity. (Owen CA Jr, Bowie EJW, Thompson JH Jr: Diagnosis of Bleeding Disorders. Second edition. Little, Brown and Company, 1975; Meijer P, Verbruggen HW, Spannagi M: Chapter 33: Clotting factors and inhibitors: Assays and Interpretation. In Laboratory Hematology Practice. Edited by K Kottke-Marchant. Wiley Blackwell Publishing, 2012, pp 435-446)

von Willebrand factor antigen, latex immunoassay:

This assay is performed using the HemosIL von Willebrand Factor Antigen kit on the Instrumentation Laboratory ACL TOP. This is a latex immunoassay method using microlatex particles coated with specific rabbit-polyclonal antibody directed against VWF. In the presence of VWF antigen, antibody-coated latex particles agglutinate to form aggregates of diameters greater than the wavelength of the light passing through the sample and more light is absorbed as aggregation increases. The increase in absorption is proportional to the concentration of VWF antigen present in the sample. (Package insert: HemosIL von Willebrand Factor Antigen, Instrumentation Laboratory, R11 05/2018)

von Willebrand factor activity, latex immunoassay:

This assay is performed using the HemosIL von Willebrand Factor Activity kit on the Instrumentation Laboratory ACL TOP. This is a latex particle-enhanced immunoassay to quantify von Willebrand factor (VWF) activity in plasma. The activity of VWF is determined by measuring the increase of turbidity produced by the agglutination of the latex reagent. A specific anti-VWF monoclonal antibody adsorbed onto the latex reagent, directed against the platelet-binding site of VWF (glycoprotein Ib receptor), reacts with the VWF of patient sample. The degree of agglutination is directly proportional to the activity of VWF in the sample and is determined by measuring the decrease of transmitted light caused by the aggregates. (Package insert: HemosIL von Willebrand Factor Activity, Instrumentation Laboratory, R7 05/2018)

PDF Report

No

Day(s) and Time(s) Test Performed

PTSC, APTSC, TTSC, CLFIB, DIMER, F8A, F_9, FXIII: Monday through Friday

VWAG, VWACT: Monday through Saturday

Analytic Time

Varies

Maximum Laboratory Time

7 days, 21 days if multimers ordered

Specimen Retention Time

7 days

Performing Laboratory Location

Rochester

Fees and Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

Test Classification

See Individual Test IDs

CPT Code Information

85610-PTSC

85730-APTSC

85670-TTSC

85384-CLFIB

85379-DIMER

85240-F8A

85250-F_9

85291-FXIII

85246-VWAG

85397-VWACT

85130-Chromogenic factor VIII (if appropriate)

85130-Chromogenic factor IX (if appropriate)

85210-Factor II (if appropriate)

85220-Factor V (if appropriate)

85230-Factor VII (if appropriate)

85245-Ristocetin cofactor (if appropriate)

85247-von Willebrand factor multimer (if appropriate)

85260-Factor X (if appropriate)

85270-Factor XI (if appropriate)

85280-Factor XII (if appropriate)

85300-Antithrombin activity (if appropriate)

85301-Antithrombin antigen (if appropriate)

85335-Bethesda units (if appropriate)

85335-Factor II inhibitor screen (if appropriate)

85335-Factor V inhibitor screen (if appropriate)

- 85335-Factor VII Inhibitor screen (if appropriate)
- 85335-Factor VIII inhibitor screen (if appropriate)
- 85335-Factor IX inhibitor screen (if appropriate)
- 85335-Factor X inhibitor screen (if appropriate)
- 85335-Factor XI inhibitor screen (if appropriate)
- 85366-Soluble fibrin monomer (if appropriate)
- 85385-PT-Fibrinogen (if appropriate)
- 85410-Alpha-2 plasmin inhibitor (if appropriate)
- 85415-PAI-1 Ag (if appropriate)
- 85420-Plasminogen Activity (if appropriate)
- 85597-Platelet neutralization for lupus inhibitor (if appropriate)
- 85598-Staclot LA (if appropriate)
- 85611-PT mix 1:1 (if appropriate)
- 85613-DRVVT (if appropriate)
- 85613-DRVVT mix (if appropriate)
- 85613-DRVVT confirm (if appropriate)
- 85635-Reptilase time (if appropriate)
- 85732-APTT mix 1:1 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
ALBLD	Bleeding Diath Prof, Limited	In Process

Result ID	Test Result Name	Result LOINC Value
CLFIB	Fibrinogen, Clauss, P	48664-7
TTSC	Thrombin Time (Bovine), P	46717-5
603322	Reviewed by	18771-6
F8A	Coag Factor VIII Activity Assay, P	3209-4
9068	Factor XIII(13),Scrn	3241-7
F_9	Coag Factor IX Assay, P	3187-2



Result ID	Test Result Name	Result LOINC Value
VWAG	von Willebrand Factor Ag, P	27816-8
APTSC	Activated Partial Thrombopl Time, P	14979-9
DIMER	D-Dimer, P	48067-3
PTSEC	Prothrombin Time (PT), P	5902-2
VWACT	von Willebrand Factor Activity, P	68324-3
INRSC	INR	6301-6
603181	Limited Bleed Prof Interp	69049-5