

Overview

Useful For

Screening for mast cell activation disorders including systemic mastocytosis using 24-hour urine specimens

Profile Information

Test Id	Reporting Name	Available Separately	Always Performed
T23BP	2,3-dinor 11B-Prostaglandin F2a	No	Yes
CRT24	Creatinine, 24 HR, U	No	Yes

Special Instructions

- [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#)

Highlights

2,3-Dinor-11beta-prostaglandin F2 alpha (2,3 BPG) is elevated in the urine of patients with systemic mastocytosis (SM).

This test should be used as a screening test for SM.

When 2,3 BPG is used in combination with urinary leukotriene E4 and N-methyl histamine, the sensitivity for SM detection increases to 90%.

Method Name

T23BP: Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

CRT24: Enzymatic Colorimetric Assay

NY State Available

Yes

Specimen

Specimen Type

Urine

Ordering Guidance

A 24-hour urine collection is the preferred specimen type, but a random specimen is also acceptable for 2,3-dinor 11beta-prostaglandin F2 alpha; order 23BPR / 2,3-Dinor 11 Beta-Prostaglandin F2 Alpha, Random, Urine.

If the total volume provided is less than 300 mL, this test will be canceled and 23BPR ordered and performed.

Necessary Information

Specimen volume in milliliters and duration are required.

Specimen Required

Patient Preparation: Patients taking aspirin or nonsteroidal anti-inflammatory drugs (NSAID) may have decreased concentrations of prostaglandin F2 alpha. If possible, discontinue for 2 weeks or 72 hours, respectively, prior to collecting a specimen.

Supplies: Sarstedt Aliquot Tube, 5 mL (T914)

Container/Tube: Plastic, 5-mL tube

Specimen Volume: 5 mL

Collection Instructions:

- 1. Collect urine for 24 hours.
- 2. No preservative preferred.

Additional Information: See [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#) for multiple collections.

Urine Preservative Collection Options

Note: The addition of preservative or application of temperature controls **must occur within 4 hours of completion** of the collection.

Ambient (no additive)	No
Refrigerate (no additive)	Preferred
Frozen (no additive)	OK
50% Acetic Acid	OK
Boric Acid	OK
Diazolidinyl Urea	No
6M Hydrochloric Acid	No
6M Nitric Acid	No
Sodium Carbonate	OK
Thymol	No
Toluene	No

Specimen Minimum Volume

2.7 mL

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
---------------	-------------	------	-------------------

Urine	Refrigerated (preferred)	14 days	
	Frozen	30 days	
	Ambient	8 hours	

Clinical & Interpretive

Clinical Information

2,3-Dinor-11beta-prostaglandin F2 alpha (2,3 BPG) is the most abundant metabolic product of prostaglandins released by activated mast cells. Systemic mastocytosis (SM) is a disease in which clonally derived mast cells accumulate in peripheral tissues. Degranulation of these mast cells releases large amounts of histamines, prostaglandins, leukotrienes, and tryptase.

World Health Organization diagnostic criteria for SM require the presence of elevated mast cell counts on a bone marrow biopsy and 1 of the following minor criteria:

- Abnormal mast cell morphology
- KIT Asp816Val variant
- CD25-positive mast cells
- Serum tryptase greater than 20 ng/mL

Alternatively, SM diagnosis can be made with the presence of 3 minor criteria in the absence of abnormal bone marrow studies.

Measurement of mast cell mediators in blood or urine is less invasive and is advised for the initial evaluation of suspected cases. Elevated levels of serum tryptase, urinary N-methylhistamine, 2,3 BPG, or leukotriene E4 are consistent with the diagnosis of systemic mast cell disease.

Reference Values

<1802 pg/mg creatinine

Interpretation

Elevated urine 2,3-dinor-11beta-prostaglandin F2 alpha is consistent with systemic mastocytosis.

Cautions

Elevated levels of 2,3-dinor-11beta-prostaglandin F2 alpha (2,3 BPG) in urine are not specific for systemic mast cell disease and may be found in patients with angioedema, diffuse urticaria, or myeloproliferative diseases in the absence of diffuse mast cell proliferation.

Systemic mast cell disease is a heterogeneous disease, and some patients may not have elevated 2,3 BPG in urine.

Clinical Reference

1. Gotlib J, Pardanani A, Akin C, et al: International Working Group-Myeloproliferative Neoplasms Research and Treatment (IWG-MRT) and European Competence Network on Mastocytosis (ECNM) consensus response criteria in advanced systemic mastocytosis. Blood. 2013 Mar 28;121(13):2393-2401. doi: 10.1182/blood-2012-09-458521

2. Butterfield JH: Increased leukotriene E4 excretion in systemic mastocytosis. Prostaglandins Other Lipid Mediat. 2010 Jun;92(1-4):73-76. doi: 10.1016/j.prostaglandins.2010.03.003

3. Roberts LJ 2nd, Sweetman BJ, Lewis RA, Austen KF, Oates JA: Increased production of prostaglandin D2 in patients with systemic mastocytosis. N Engl J Med. 1980 Dec 11;303(24):1400-1404. doi: 10.1056/NEJM198012113032405

4. Metcalfe DD: Mastocytosis syndromes. In: Middleton E Jr, Reed CE, Ellis EF, et al. eds. Allergy Principles and Practice. Vol II. 4th ed. Mosby Yearbook Inc; 1993:1537-1551

5. Butterfield J, Weiler CR: The utility of measuring urinary metabolites of mast cell mediators in systemic mastocytosis and mast cell activation syndrome. J Allergy Clin Immunol Pract. 2020 Sept;8(8):2533-2541

Performance

Method Description

2,3-Dinor-11beta-prostaglandin F2 alpha (2,3 BPG) is quantified in urine by liquid chromatography tandem mass spectrometry.(Unpublished Mayo method)

Creatinine:

All 2,3 BPG concentrations are normalized to urine creatinine levels measured using a Roche cobas enzymatic method. The enzymatic method is based on the determination of sarcosine from creatinine with the aid of creatininase, creatinase, and sarcosine oxidase. The liberated hydrogen peroxide is measured via a modified Trinder reaction using a colorimetric indicator. Optimization of the buffer system and the colorimetric indicator enables the creatinine concentration to be quantified both precisely and specifically.(Package insert: Creatinine plus ver 2. Roche Diagnostics; V15.0, 03/2019)

PDF Report

No

Day(s) Performed

Tuesday, Thursday

Report Available

3 to 8 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

84150

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
23BPT	2,3-dinor 11B-Prostaglandin F2a, U	94381-1

Result ID	Test Result Name	Result LOINC® Value
TM10	Collection Duration	13362-9
VL8	Urine Volume	3167-4
CR_A	Creatinine, 24 HR, U	2162-6
CR_24	Creatinine Concentration, 24 HR, U	20624-3
603460	2,3-dinor 11B-Prostaglandin F2a	94381-1