

Overview

Useful For

Detecting drug abuse involving barbiturates such as amobarbital, butalbital, pentobarbital, phenobarbital, and secobarbital

Chain of custody is required whenever the results of testing could be used in a court of law. Its purpose is to protect the rights of the individual contributing the specimen by demonstrating that, at all times, it was under the control of personnel involved with testing the specimen; this control implies that the opportunity for specimen tampering would be limited.

Additional Tests

Test Id	Reporting Name	Available Separately	Always Performed
COCH	Chain of Custody Processing	No	Yes
ADLTX	Adulterants Survey, CoC, U	Yes	Yes

Testing Algorithm

Adulterants testing will be performed on all chain-of-custody urine samples as per regulatory requirements.

Method Name

BARBX: Immunoassay/Gas Chromatography Mass Spectrometry (GC-MS) Confirmation with Quantitation  
ADLTX: Spectrophotometry (SP)

NY State Available

Yes

Specimen

Specimen Type

Urine

Specimen Required

**Supplies:** Chain of Custody Kit (T282)  
**Container/Tube:** Chain of custody kit (T282) containing the specimen containers, seals, and documentation required  
**Specimen Volume:** 10 mL  
**Collection Instructions:** Collect specimen in the container provided, seal, and submit with the associated documentation to satisfy the legal requirements for chain-of-custody testing.  
**Additional Information:** Submitting less than 10 mL will compromise the ability to perform all necessary testing.

Forms

1. [Chain of Custody Request](#) is included in the Chain of Custody Kit (T282).
2. If not ordering electronically, complete, print, and send a [Therapeutics Test Request](#) (T831) with the specimen.

Specimen Minimum Volume

5 mL

Reject Due To

Gross hemolysis	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	14 days	
	Frozen	14 days	
	Ambient	72 hours	

Clinical & Interpretive

Clinical Information

Barbiturates represent a class of drugs that were originally introduced as sleep inducers. Butalbital is also used to control severe headaches. Mephobarbital and phenobarbital are frequently used to control major motor (grand mal) seizures. These drugs are commonly abused as "downers" to induce sleep after an amphetamine- or cocaine-induced "high."

Chain of custody is a record of the disposition of a specimen to document the personnel who collected, handled, and performed the analysis. When a specimen is submitted in this manner, analysis will be performed in such a way that it will withstand regular court scrutiny.

Reference Values

Negative

Cutoff concentrations:

IMMUNOASSAY SCREEN

<200 ng/mL

GAS CHROMATOGRAPHY MASS SPECTROSCOPY:

Butalbital: <100 ng/mL

Amobarbital <100 ng/mL

Pentobarbital <100 ng/mL

Secobarbital <100 ng/mL  
Phenobarbital <100 ng/mL

**Interpretation**

The presence of a barbiturate in urine indicates use of one of these drugs.

Most of the barbiturates are fast acting; their presence indicates use within the past 3 days.

Phenobarbital, commonly used to control epilepsy, has a very long half-life. The presence of phenobarbital in urine indicates that the patient has used the drug sometime within the past 30 days.

**Cautions**

No significant cautionary statements

**Clinical Reference**

1. Baselt RC, Cravey RH. Disposition of Toxic Drugs and Chemicals in Man. 3rd ed. Year Book Medical Publishers; 1989
2. Langman LJ, Bechtel LK, Meier BM, Holstege C. Clinical toxicology. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:832-887

**Performance****Method Description**

Preliminary screen is performed by immunoassay.

The barbiturate assay is based on the kinetic interaction of microparticles in a solution as measured by changes in light transmission. In the absence of sample drug, soluble drug conjugates bind to antibody-bound microparticles causing the formation of particle aggregates. As the aggregation reaction proceeds in the absence of sample drug, the absorbance increases. When a urine sample contains the drug in question, this drug competes with the drug derivative conjugate for microparticle-bound antibody. Antibody bound to sample drug is no longer available to promote particle aggregation, and subsequent particle lattice formation is inhibited. The presence of sample drug diminishes the increasing absorbance in proportion to the concentration of drug in the sample. Sample drug content is determined relative to the value obtained for a known cutoff concentration of drug.(Package insert: Barbiturates. Roche Diagnostics; 03/2019)

The specimen is then analyzed by gas chromatography mass spectrometry (GC/MS) for confirmation with quantitation.(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

Tuesday, Thursday, Sunday

**Report Available**

3 to 5 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80345

G0480 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
BARBX	Barbiturates Confirmation, CoC, U	53746-4

Result ID	Test Result Name	Result LOINC® Value
6668	Barbiturates Immunoassay Screen	70155-7
36143	Butalbital-by GC/MS	16237-0
36144	Amobarbital-by GC/MS	16239-6
36145	Pentobarbital-by GC/MS	16240-4
36146	Secobarbital-by GC/MS	16238-8
36147	Phenobarbital-by GC/MS	16241-2
36148	Barbiturates Interpretation	69050-3
36149	Chain of Custody	77202-0