

### Leukocyte Cell-Derived Chemotaxin 2 (LECT2), Immunostains Without Interpretation

#### Overview

#### Useful For

Identification and classification of amyloid subtypes in tissue

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

#### **Testing Algorithm**

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

#### Method Name

Immunohistochemistry (IHC)

#### NY State Available

Yes

#### Specimen

Specimen Type

TECHONLY

#### **Ordering Guidance**

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

#### **Shipping Instructions**

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

#### **Specimen Required**

Supplies: Immunostain Technical Only Envelope (T693)
Specimen Type: Tissue
Container/Tube: Immunostains Technical Only Envelope
Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick.



### Leukocyte Cell-Derived Chemotaxin 2 (LECT2), Immunostains Without Interpretation

Acceptable: Formalin-fixed, paraffin-embedded tissue block

#### Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <u>https://news.mayocliniclabs.com/pathology/digital-imaging/</u>

2. Clients ordering stains using a manual requisition form will not have access to digital images.

3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <u>https://news.mayocliniclabs.com/pathology/digital-imaging/#section3</u>

#### Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u> <u>Request</u> (T763) with the specimen.

#### Reject Due To

Wet/frozen	Reject
tissue Cytology	
smears	
Nonformalin	
fixed tissue	
Nonparaffin	
embedded	
tissue	
Noncharged	
slides ProbeOn	
slides	

#### **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

#### **Clinical & Interpretive**

#### **Clinical Information**

Immunohistochemical staining for leukocyte cell-derived chemotaxin 2 (LECT2) is useful in the process of confirming amyloid subtype. Antibodies to LECT2 stain the amyloid deposits in patients with LECT2 amyloidosis. LECT2 amyloidosis typically involves the kidney, liver, and spleen.

#### Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.



Leukocyte Cell-Derived Chemotaxin 2 (LECT2), Immunostains Without Interpretation

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

#### Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

#### **Clinical Reference**

<u>1</u>1. Benson MD, James S, Scott K, et al. Leukocyte chemotactic factor 2: A novel renal amyloid protein. Kidney Int. 2008;74(2):218-222

2. Uchida T, Nagai H, Gotoh K, et al. Expression pattern of a newly recognized protein, LECT2, in hepatocellular carcinoma and its premalignant lesion. Pathol Int. 1999;49(2):147-151

3. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5\_25

#### Performance

#### Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed Monday through Friday

Report Available 1 to 3 days

**Specimen Retention Time** Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes



# Leukocyte Cell-Derived Chemotaxin 2 (LECT2),

### Immunostains Without Interpretation

#### Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information. ٠
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week. ٠
- Prospective clients should contact their account representative. For assistance, contact Customer Service. •

#### **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

#### **CPT Code Information**

88342-TC, primary 88341-TC, if additional IHC

#### LOINC<sup>®</sup> Information

Test ID	Test Order Name	Order LOINC <sup>®</sup> Value
LECT2	LECT2 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC <sup>®</sup> Value
70809	LECT2 IHC, Tech Only	Bill only; no result