

S-100 Immunostain, Technical Component Only

#### Overview

#### Useful For

Aids in the identification of various neoplasms

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

#### **Testing Algorithm**

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

#### Method Name

Immunohistochemistry (IHC)

#### NY State Available

Yes

#### Specimen

## Specimen Type

TECHONLY

#### **Ordering Guidance**

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

#### **Shipping Instructions**

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

#### **Specimen Required**

Supplies: Immunostain Technical Only Envelope (T693)
Specimen Type: Tissue
Container/Tube: Immunostain Technical Only Envelope
Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick.



S-100 Immunostain, Technical Component Only

Acceptable: Formalin-fixed, paraffin-embedded tissue block

#### Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <a href="https://news.mayocliniclabs.com/pathology/digital-imaging/">https://news.mayocliniclabs.com/pathology/digital-imaging/</a>

2. Clients ordering stains using a manual requisition form will not have access to digital images.

3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <u>https://news.mayocliniclabs.com/pathology/digital-imaging/#section3</u>

#### Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u> <u>Request</u> (T763) with the specimen.

### Reject Due To

Wet/frozen	Reject
tissue	
Cytology	
smears	
Nonformalin	
fixed tissue	
Nonparaffin	
embedded	
tissue	
Noncharged	
slides	
ProbeOn slides	

#### **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

### **Clinical & Interpretive**

#### **Clinical Information**

S100 protein expression is seen in cartilaginous tumors, myoepithelial tumors, Schwann cells and neural tumors, Langerhans cell proliferations, benign and malignant melanocytes, clear cell sarcoma, and some carcinomas (particularly of the breast). S100 staining occurs both in the nucleus and cytoplasm.

#### Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is



S-100 Immunostain, Technical Component Only

required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

#### Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

#### **Clinical Reference**

1. Blessing K, Sanders DS, Grant JJ. Comparison of immunohistochemical staining of the novel antibody melan-A with S100 protein and HMB-45 in malignant melanoma and melanoma variants. Histopathology. 1998;32(2):139-146. doi:10.1046/j.1365-2559.1998.00312.x

2. Clarkson KS, Sturdgess IC, Molyneux AJ. The usefulness of tyrosinase in the immunohistochemical assessment of melanocytic lesions: a comparison of the novel T311 antibody (anti-tyrosinase) with S-100, HMB45, and A103 (anti-melan-A). J Clin Pathol. 2001;54(3):196-200

 Matsunou H, Shimoda T, Kakimoto S, Yamashita H, Ishikawa E, Mukai M. Histopathologic and immunohistochemical study of malignant tumors of peripheral nerve sheath (malignant schwannoma). Cancer. 1985;56(9):2269-2279
Orchard GE. Comparison of immunohistochemical labelling of melanocyte differentiation antibodies melan-A, tyrosinase and HMB 45 with NKIC3 and S100 protein in the evaluation of benign naevi and malignant melanoma. Histochem J. 2000;32(8):475-481. doi:10.1023/a:1004192232357

5. Al-Ismaeel Q, Neal CP, Al-Mahmoodi H, et al. ZEB1 and IL-6/11-STAT3 signalling cooperate to define invasive potential of pancreatic cancer cells via differential regulation of the expression of S100 proteins. Br J Cancer. 2019;121(1):65-75. doi:10.1038/s41416-019-0483-9

6. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5\_25

### Performance

#### **Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report No

Day(s) Performed Monday through Friday

**Report Available** 

1 to 3 days



S-100 Immunostain, Technical Component Only

#### **Specimen Retention Time**

Until staining is complete.

#### **Performing Laboratory Location**

Rochester

## Fees & Codes

#### Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

#### **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

#### **CPT Code Information**

88342-TC, primary 88341-TC, if additional IHC

### LOINC<sup>®</sup> Information

Test ID	Test Order Name	Order LOINC <sup>®</sup> Value
S100	S-100 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC <sup>®</sup> Value
70859	S-100 IHC, Tech Only	Bill only; no result