

Overview**Method Name**

Double Immunodiffusion (DID)

NY State Available

Yes

Specimen**Specimen Type**

Serum

Specimen Required**Specimen Type:** Serum**Container/Tube:** Red top or SST**Specimen Volume:** 1 mL**Collection Instructions:** Draw blood in a plain red-top tube, serum gel tube(s) is also acceptable. Separate serum immediately after coagulation (30 minutes) to prevent hemolysis. Send 1 mL of serum frozen in a plastic vial.

NOTE: Patient should be fasting for eight hours to avoid lipemic sample interference.

Specimen Minimum Volume

0.5 mL

Reject Due To

Hemolysis	Mild reject; Gross reject
Lipemia	Mild reject; Gross reject
Icterus	NA
Other	Gross bacterial contamination

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Frozen (preferred)	14 days	
	Refrigerated	72 hours	

Clinical and Interpretive

Reference Values

Aspergillus fumigatus Neg: <1:1

Aspergillus flavus Neg: <1:1

Aspergillus niger Neg: <1:1

Performance**PDF Report**

No

Day(s) and Time(s) Test Performed

Monday through Friday

Analytic Time

3-5 days

Maximum Laboratory Time

5-9 days

Performing Laboratory Location

LabCorp Burlington

Fees and Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

CPT Code Information

86606 x 3

LOINC® Information

Test ID	Test Order Name	Order LOINC Value
FASAB	Aspergillus Ab, Qn, DID	Not Provided

Result ID	Test Result Name	Result LOINC Value
Z5651	Aspergillus fumigatus	23821-2
Z5652	Aspergillus flavus	23820-4
Z5653	Aspergillus niger	10894-4