

Overview

Method Name

RIPA Gel Radiography

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Collection Container/Tube: Red

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Acceptable: 5 mL SST

Collection Instructions: Draw blood in a plain red-top tube(s), serum gel tube is acceptable. Spin down and send 1 mL of serum refrigerated in a plastic vial.

Specimen Minimum Volume

0.3 mL (volume does NOT allow for repeat testing)

Reject Due To

Gross hemolysis:	Reject
Thawing:	Warm OK; Cold OK
Gross lipemia:	Reject
Gross icterus	Reject
Other:	Anything other than serum; bacterial contamination

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	60 days	
	Ambient	7 days	

Clinical & Interpretive

Clinical Information

The Th/To antibodies are present in 10-19% of patients with limited SSc, in 11% of patients with diffuse cutaneous SSc, and in 3% of patients with primary Raynaud's disease. Anti-Th/To antibody has been shown to be highly specific for patients with SSc.

Reference Values

Negative

Performance

PDF Report

No

Report Available

16-25 days

Performing Laboratory Location

Esoterix Endocrinology

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

CPT Code Information

83516

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FATHO	Anti-Th/To Ab	81743-7

Result ID	Test Result Name	Result LOINC® Value
-----------	------------------	---------------------

FATHO	Anti-Th/To Ab	81743-7
-------	---------------	---------