

Overview

Useful For

Monitoring secobarbital therapy

Method Name

Gas Chromatography Mass Spectrometry (GC-MS)

NY State Available

Yes

Specimen

Specimen Type

Serum Red

Specimen Required

Collection Container/Tube: Red top (Serum gel/SST are **not acceptable**)

Submission Container/Tube: Plastic vial

Specimen Volume: 1.2 mL

Collection Instructions:

1. Draw blood immediately before next scheduled dose.
2. Centrifuge and aliquot serum in plastic vial within 2 hours of collection.

Forms

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

-[Neurology Specialty Testing Client Test Request](#) (T732)

-[Therapeutics Test Request](#) (T831)

Specimen Minimum Volume

0.6 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	14 days	
	Ambient	14 days	
	Frozen	14 days	

Clinical & Interpretive

Clinical Information

Secobarbital is a short-acting barbiturate with hypnotic properties used as a preanesthetic agent and in the short-term treatment of insomnia.(1,2)

Secobarbital is administered orally. The duration of its hypnotic effect is about 3 to 4 hours. The drug distributes throughout the body, with a volume of distribution of 1.6 to 1.9 L/kg, and about 46% to 70% of a dose is bound to plasma proteins. Metabolism takes place in the liver primarily via hepatic microsomal enzymes. The half-life of secobarbital is about 15 to 40 hours (mean: 28 hours).(2,3)

Reference Values

Therapeutic concentration: 1.0-2.0 mcg/mL

Toxic concentration: >5.0 mcg/mL

Interpretation

Secobarbital concentrations above 5 mcg/mL have been associated with toxicity.

Cautions

The concentration at which toxicity occurs varies and results should be interpreted in light of the clinical situation.

Specimens collected in serum gel tubes are not acceptable because the drug can absorb on the gel and lead to falsely decreased concentrations.

Clinical Reference

1. Mihic SJ, Mayfield J, Harris RA: Hypnotics and sedatives. In: Brunton LL, Hilal-Dandan R, Knollmann BC, eds. Goodman and Gilman's The Pharmacological Basis of Therapeutics. 13th ed. McGraw-Hill Education; 2017

2. Milone MC, Shaw LM: Therapeutic drugs and their management. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:800-831

3. Baselt RC: Disposition of Toxic Drugs and Chemicals in Man. 10th ed. Biomedical Publications; 2014:2211

4. Langman LJ, Bechtel LK, Meier BM, Holstege C: Clinical toxicology. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. Elsevier; 2018:832-887

Performance

Method Description

Barbiturates are extracted from serum using solid-phase extraction techniques. The serum is buffered and eluted with organic solvent. The organic phase is dried, reconstituted, and analysis performed by gas chromatography-mass

spectrometry using selected ion monitoring. The assay utilizes deuterated barbiturates as internal standards.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Wednesday

Report Available

3 to 9 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80345
G0480 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
SECOS	Secobarbital, S	4028-7

Result ID	Test Result Name	Result LOINC® Value
8243	Secobarbital, S	4028-7