

## Overview

### Useful For

Evaluation of a prolonged thrombin time (TT) (mainly used to confirm or exclude the presence of heparin in the specimen)

Evaluating hypofibrinogenemia or dysfibrinogenemia in conjunction with the TT and fibrinogen assay

### Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)

### Method Name

Optical Clot-Based

### NY State Available

Yes

## Specimen

### Specimen Type

Plasma Na Cit

### Specimen Required

**Specimen Type:** Platelet-poor plasma

**Collection Container/Tube:** Light-blue top (citrate)

**Submission Container/Tube:** Plastic vial

**Specimen Volume:** 1 mL Platelet poor plasma

#### Collection Instructions:

1. For complete instruction, see [Coagulation Guidelines for Specimen Handling and Processing](#).
2. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
3. Immediately freeze plasma (no longer than 4 hours after collection) at -20 degrees C or, ideally at -40 degrees C or below.

#### Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

### Forms

If not ordering electronically, complete, print, and send a [Coagulation Test Request](#) (T753) with the specimen.

### Specimen Minimum Volume

Platelet poor plasma: 0.5 mL

## Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

## Clinical & Interpretive

### Clinical Information

Prolonged clotting times may be associated with a wide variety of coagulation abnormalities including:

- Deficiency or functional abnormality (congenital or acquired) of any of the coagulation proteins
- Deficiency or functional abnormality of platelets
- Specific factor inhibitors
- Acute disseminated intravascular coagulation
- Exogenous anticoagulants (eg, heparin, warfarin)

The prothrombin time (PT) and activated partial thromboplastin time (APTT) are first-order tests for coagulation abnormalities and are prolonged in many bleeding disorders. A battery of coagulation tests is often required to determine the cause of prolonged clotting times. The thrombin time (TT) test is used to identify the cause of prolonged APTT or dilute Russell's viper venom time (DRVVT). Reptilase time (RT) test is used to evaluate a prolonged TT.

Reptilase is a thrombin-like enzyme isolated from the venom of *Bothrops atrox*. Thrombin splits small fibrinopeptides A and B from fibrinogen molecules, producing fibrin monomer, which polymerizes to form a clot. Reptilase, however, splits off fibrinopeptide A but not B, which results in fibrin polymerization. In contrast to thrombin and the TT test which are inhibited by heparin, the RT is normal in the presence of heparin. Similar to the TT test, the RT is prolonged in the presence of hypofibrinogenemia and dysfibrinogenemia.

### Reference Values

14.0-23.9 Seconds

### Interpretation

As seen in the following table, reptilase time can help distinguish among the various causes of a prolonged thrombin time (TT).

Table.

Thrombin time	Reptilase time	Causes	Remarks

Prolonged	Prolonged	Hypo- or afibrinogenemia	Ascertain by determination of fibrinogen
Prolonged	Prolonged	Dysfibrinogenemia	Ascertain by specific assay
Prolonged	Normal	Heparin or inhibitor of thrombin	Differentiate by human TT and/or heparin assays
Prolonged	Prolonged	Fibrin(ogen) split products (FSP)	Ascertain by FSP or D-dimer assay

## Cautions

The reptilase time test has limited diagnostic value when ordered as a stand-alone test.

## Clinical Reference

Favaloro EJ, Lippi G, eds. Hemostasis and Thrombosis: Methods and Protocols. Humana Press; 2017

## Performance

### Method Description

The reptilase time assay is performed on the Instrumentation Laboratories ACL TOP. Patient plasma is combined with a reptilase reagent containing a thrombin-like enzyme triggering the coagulation process in the mixture. Time to clot formation is measured optically using a wavelength of 671 nm. (Owen CA, Bowie EJW, Thompson JH. Tests of hemostasis and blood coagulation. In: The Diagnosis of Bleeding Disorders. 2nd ed. Little, Brown, and Company; 1975:85-154; package insert: STA-Reptilase. Diagnostica Stago, Inc; 03/2018)

### PDF Report

No

### Day(s) Performed

Monday through Friday

### Report Available

1 to 4 days

### Specimen Retention Time

7 days

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.

- 
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

85635

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
RTSC	Reptilase Time, P	6683-7

Result ID	Test Result Name	Result LOINC® Value
RTSC	Reptilase Time, P	6683-7