



Test Definition: SP142

Programmed Death-Ligand 1 (PD-L1) (SP142),
Semi-Quantitative Immunohistochemistry,
Manual

Overview

Useful For

Identification of neoplasms expressing programmed cell death 1-ligand 1 (clone SP142)

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

Special

Ordering Guidance

[In patients with specific tumor types, programmed death-ligand 1 \(PD-L1\) immunohistochemistry \(IHC\) is indicated to predict response to treatment with PD-L1 inhibitors. The specific PD-L1 clone, scoring method, and eligibility requirements depend on the tumor type, stage of malignancy, previous treatment outcomes, and specific PD-L1 inhibitor under consideration. For assistance with PD-L1 test selection as well as answers to frequently asked questions, see \[PD-L1 Immunohistochemistry Testing\]\(#\) on MayoClinicLabs.com.](#)

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) to the outside of the transport container before putting into the courier mailer.

Necessary Information

A pathology/diagnostic report and a brief history, including primary site of neoplasm, are required.

Specimen Required

This assay requires at least 100 viable tumor cells.

Specimen Type: Tissue

Supplies: Pathology Packaging Kit (T554)

Submit:

Formalin-fixed, paraffin-embedded tissue block

OR

3 Unstained glass, "positively charged" slides with 4-microns formalin-fixed, paraffin-embedded tissue

Additional Information: One slide will be stained with hematoxylin and eosin and returned.

Forms

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

[Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request \(T763\)](#)

[Oncology Test Request \(T729\)](#)

Reject Due To

| | |
|---|--------|
| Decalcified paraffin embedded tissue | Reject |
| Wet/frozen tissue | Reject |
| Cytology smears | Reject |
| Nonformalin fixed tissue including alcohol-formalin-acetic acid (AFA), 95% ethanol, PREFER fixatives or zinc formalin | Reject |
| Nonparaffin embedded tissue | Reject |
| Noncharged slides | Reject |
| ProbeOn slides | Reject |
| Snowcoat slides | Reject |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|---------------------|------|-------------------|
| Special | Ambient (preferred) | | |
| | Refrigerated | | |

Clinical & Interpretive**Clinical Information**

Programmed cell death 1-ligand 1 (PD-L1), also known as B7 homolog 1 (B7-H1) or CD274, is a transmembrane protein involved in the regulation of cell-mediated immune responses through interaction with the receptor programmed death protein-1. PD-L1 has been identified as both a prognostic and therapeutic marker in a variety of neoplasms. Overexpression of PD-L1 has been observed in carcinomas of the urinary bladder, lung, thymus, colon, pancreas, ovary, breast, kidney, and in melanoma and glioblastoma.

Interpretation

The results of the test will be reported in the form of scores. The scoring system is based on type and origin of tumor. If additional interpretation or analysis is needed, order PATHC / Pathology Consultation along with this test.

Cautions

Preclinical studies suggest that positive programmed cell death 1-ligand 1 (PD-L1) immunohistochemistry in tumor cells may predict tumor response to therapy with immune checkpoint inhibitors. This result should not be used as the sole factor in determining treatment, as other factors (eg, tumor mutation burden and microsatellite instability) have also been studied as predictive markers.

This test has been validated for non-decalcified paraffin-embedded tissue specimens fixed in 10% neutral buffered formalin at Mayo Clinic in Rochester, Minnesota. Specimens are recommended to be placed in formalin within 1 hour of acquisition and fixed between 6 and 72 hours. This assay has not been validated on tissue or cellblocks subjected to alternative fixatives or decalcification.

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- Minimize time slides are stored after being unpackaged
- Limit exposure to high humidity and heat
- Minimize exposure to plastics

Clinical Reference

1. Rimm DL, Han G, Taube JM, et al. A prospective, multi-institutional, pathologist-based assessment of 4 immunohistochemistry assay for PD-L1 expression in non-small cell lung cancer. *JAMA Oncol.* 2017;3(8):1051-1058. doi:10.1001/jamaoncol.2017.0013
2. Gaule P, Smithy JW, Toki M, et al. A quantitative comparison of antibodies to programmed cell death 1 Ligand 1. *JAMA Oncol.* 2017;3(2):256-259. doi:10.1001/jamaoncol.2016.3015
3. Sunshine JC, Nguyen P, Kaunitz G, et al. PD-L1 expression in melanoma: A quantitative immunohistochemical antibody

comparison. Clin Can Res. 2017;23(16):4938-4944. doi:10.1158/1078-0432.CCR-16-1821

4. D'Incecco A, Andrezzi M, Ludovini V, et al. PD-1 and PD-L1 expression in molecularly selected non-small-cell lung cancer patients. Br J Cancer. 2015;112(1):95-102. doi:10.1038/bjc.2014.555

5. Mansfield AS, Roden AC, Peikert T, et al. B7-H1 expression in malignant pleural mesothelioma is associated with sarcomatoid histology and poor prognosis. J Thorac Oncol. 2014;9(7):1036-1040. doi:10.1097/JTO.000000000000177

6. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue using Ventana PD-L1 clone SP142.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

5 to 7 days

Specimen Retention Time

Until reported

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88360

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|--|--------------------|
| SP142 | PD-L1 (SP142) SemiQuant IHC, Manual | 85149-3 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|------------------------------------|---------------------|
| 603770 | Interpretation | 59465-5 |
| 603771 | Participated in the Interpretation | No LOINC Needed |
| 603772 | Report electronically signed by | 19139-5 |
| 603773 | Material Received | 81178-6 |
| 603774 | Disclaimer | 62364-5 |
| 603775 | Case Number | 80398-1 |