

Overview

Useful For

Managing everolimus immunosuppression in solid organ transplant

Method Name

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

NY State Available

No

Specimen

Specimen Type

Whole Blood EDTA

Specimen Required

Container/Tube: Lavender top (EDTA)

Specimen Volume: 3 mL

Collection Instructions:

1. Draw blood immediately before next scheduled dose.
2. **Do not centrifuge.**
3. Send whole blood specimen in original tube. **Do not aliquot.**

Additional Information: Therapeutic range applies to trough specimens collected immediately prior to a.m. dose.

Forms

If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:

[-Renal Diagnostics Test Request \(T830\)](#)

[-Therapeutics Test Request \(T831\)](#)

Specimen Minimum Volume

1 mL

Reject Due To

| | |
|-------------------|--------|
| Gross hemolysis | OK |
| Gross lipemia | OK |
| Gross icterus | OK |
| Clotted specimens | Reject |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|------------------|--------------------------|---------|-------------------|
| Whole Blood EDTA | Refrigerated (preferred) | 28 days | |
| | Frozen | 28 days | |
| | Ambient | 14 days | |

Clinical & Interpretive**Clinical Information**

Everolimus is an immunosuppressive agent derived from sirolimus (rapamycin). Both drugs function via inhibition of mTOR (mechanistic target of rapamycin) signaling and share similar pharmacokinetic and toxicity profiles. Everolimus has a shorter half-life than sirolimus, which allows for more rapid achievement of steady-state pharmacokinetics. Everolimus is extensively metabolized, primarily by cytochrome P450 (CYP) 3A4, thus its use with inducers or inhibitors of that enzyme may require dose adjustment. The most common adverse effects include hyperlipidemia, thrombocytopenia, and nephrotoxicity. Everolimus is useful as adjuvant therapy in renal cell carcinoma and other cancers. It recently gained US Food and Drug Administration approval for prophylaxis of graft rejection in solid organ transplant, an application that has been accepted for years in Europe. The utility of therapeutic drug monitoring has not been established for everolimus as an oncology chemotherapy agent; however, measuring blood drug concentrations is common practice for its use in transplant. Therapeutic targets vary depending on the transplant site and institution protocol. Guidelines for heart and kidney transplants suggest that trough (immediately prior to the next scheduled dose) blood concentrations between 3 and 8 ng/mL provide optimal outcomes.

Reference Values

3-8 ng/mL

Target steady-state trough concentrations vary depending on the type of transplant, concomitant immunosuppression, clinical/institutional protocols, and time post-transplant. Results should be interpreted in conjunction with this clinical information and any physical signs/symptoms of rejection/toxicity.

Interpretation

Therapeutic targets vary by transplant site and institution protocol. Heart and kidney transplant guidelines suggest a therapeutic range of 3 to 8 ng/mL.

Measurement of drug concentrations in oncology chemotherapy is less common, thus no therapeutic range is established for this application.

Cautions

Therapeutic targets vary by transplant site and institution protocol. Established ranges refer to trough (predose) concentrations.

Clinical Reference

1. Eisen HJ, Tuzcu EM, Dorent R, et al: Everolimus for the prevention of allograft rejection and vasculopathy in cardiac-transplant recipients. *N Engl J Med.* 2003;349(9):847-858
2. Kovarik JM, Beyer D, Schmouder RL: Everolimus drug interactions: application of a classification system for clinical

decision making. *Biopharm Drug Dispos.* 2006;27(9):421-426

3. Rothenburger M, Zuckermann A, Bara C, et al: Recommendations for the use of everolimus (Certican) in heart transplantation: results from the second German-Austrian Certican Consensus Conference. *J Heart Lung Transplant.* 2007;26(4):305-311

4. Sanchez-Fructuoso AI: Everolimus: an update on the mechanism of action, pharmacokinetics and recent clinical trials. *Expert Opin Drug Metab Toxicol.* 2008;4(6):807-819

5. [Milone MC, Shaw LM: Therapeutic drugs and their management.](#) In: Rifai N, Horvath AR, Wittwer CT, eds. *Tietz Textbook of Clinical Chemistry and Molecular Diagnostics.* 6th ed. Elsevier; 2018:800-831

6. van Gelder T, Fischer L, Shihab F, Shipkova M: Optimizing everolimus exposure when combined with calcineurin inhibitors in solid organ transplantation. *Transplant Rev (Orlando).* 2017;31(3):151-157. doi:10.1016/j.trre.2017.02.007

Performance

Method Description

Whole blood samples are mixed with methanolic zinc sulfate to lyse blood cells. The supernatant is removed and analyzed by liquid chromatography-tandem mass spectrometry. (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

Same day/1 to 2 days

Specimen Retention Time

14 days

Performing Laboratory Location

Jacksonville

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

80169

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|-----------------|--------------------|
| EVROL | Everolimus, B | 50544-6 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|------------------|---------------------|
| 35146 | Everolimus, B | 50544-6 |