

## Overview

### Useful For

Assessing the cause of abnormal serum magnesium concentrations using a 24-hour urine collection

Determining whether nutritional magnesium loads are adequate

Calculating urinary calcium oxalate and calcium phosphate supersaturation and assessing kidney stone risk

### Special Instructions

- [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#)

### Method Name

Colorimetric Endpoint Assay

### NY State Available

Yes

## Specimen

### Specimen Type

Urine

### Necessary Information

**Specimen volume in milliliters and duration are required.**

### Specimen Required

**Supplies:** Sarstedt 5 mL Aliquot Tube (T914)

**Collection Container/Tube:** 24-hour graduated urine container with no metal cap or glued insert

**Submission Container/Tube:** Plastic, 5 mL tube or a clean, plastic urine container with no metal cap or glued insert

**Specimen Volume:** 4 mL

#### Collection Instructions:

1. Collect urine for 24 hours.
2. Refrigerate specimen within 4 hours of completion of 24-hour collection.

**Additional Information:** See [Urine Preservatives-Collection and Transportation for 24-Hour Urine Specimens](#) for multiple collections.

### Forms

If not ordering electronically, complete, print, and send a [Renal Diagnostics Test Request](#) (T830) with the specimen.

### Urine Preservative Collection Options

**Note:** The addition of preservative or application of temperature controls **must occur within 4 hours of completion** of the collection.

Ambient	OK
Refrigerate	Preferred
Frozen	OK
50% Acetic Acid	OK
Boric Acid	OK
Diazolidinyl Urea	OK
6M Hydrochloric Acid	OK
6M Nitric Acid	No
Sodium Carbonate	No
Toluene	No

## Specimen Minimum Volume

1 mL

## Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Urine	Refrigerated (preferred)	14 days	
	Ambient	72 hours	
	Frozen	30 days	

## Clinical & Interpretive

### Clinical Information

Magnesium, along with potassium, is a major intracellular cation. Magnesium is a cofactor of many enzyme systems. All adenosine triphosphate-dependent enzymatic reactions require magnesium as a cofactor. Approximately 70% of magnesium ions are stored in bone. The remainder are involved in intermediary metabolic processes; about 70% are present in free form, while the other 30% are bound to proteins (especially albumin), citrates, phosphate, and other complex formers. The serum magnesium level is kept constant within very narrow limits

Renal handling of magnesium is determined by the combination of filtration and reabsorption. Roughly 70% of the magnesium in plasma is filtered by the glomeruli; 20% to 30% of the filtered magnesium is reabsorbed in the proximal tubule, while less than 5% is reabsorbed in the distal tubule and collecting duct.(1)

Numerous causes of renal magnesium wasting have been identified including (but not limited to) congenital defects (including Barter and Gitelman syndrome), various endocrine disorders (including hyperaldosteronism and hyperparathyroidism), exposure to certain drugs (ie, diuretics, cis-platinum, aminoglycoside antibiotics, calcineurin

inhibitors), and other miscellaneous causes (including chronic alcohol abuse). Gastrointestinal conditions associated with fat malabsorption and chronic diarrhea can cause fecal magnesium loss and hypomagnesemia. High levels of plasma magnesium are typically only seen in patients with decreased renal function, after administration of a magnesium load large enough to exceed the kidneys' ability to excrete it, or a combination of the 2.(2)

Magnesium is an inhibitor of calcium crystal growth and contributes to urinary calcium oxalate and calcium phosphate supersaturation. However, low urinary magnesium in isolation has not been identified as a common cause of kidney stones, nor has magnesium supplementation been proven as an effective therapy for stone prevention.

**Reference Values**

51-269 mg/24 hours

Reference values have not been established for patients <18 years and >83 years of age.

**Interpretation**

Urinary magnesium excretion should be interpreted in concert with serum concentrations.

In the presence of hypomagnesemia, a 24-hour urine magnesium greater than 24 mg/day or fractional excretion greater than 0.5% suggests renal magnesium wasting. Lower values suggest inadequate magnesium intake and/or gastrointestinal losses.

In the presence of hypermagnesemia, urinary magnesium levels provide an indication of current magnesium intake.

Lower urinary magnesium excretion increases urinary calcium oxalate and calcium phosphate supersaturation and could contribute to kidney stone risk.

**Cautions**

Urinary magnesium excretion must be interpreted with caution during periods of intravenous magnesium infusion.

**Clinical Reference**

1. Delaney MP, Lamb EJ: Kidney disease. In: Rifai N, Horvath AR, Wittwer CT, eds. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. 6th ed. 2018:1309
2. Al Ghamdi SM: Magnesium deficiency: pathophysiologic and clinical overview. Am J Kidney Dis. 1994;24(5):737-752
3. Sutton RA: Abnormal renal magnesium handling. Miner Electrolyte Metab. 1993;19(4-5):232-240

**Performance****Method Description**

In alkaline solution, magnesium forms a purple complex with xylydyl blue, diazonium salt. The magnesium concentration is measured photometrically via the decrease in xylydyl blue absorbance.(Package insert: Roche MG2 kit. Roche Diagnostics; V8.0 01/2020)

**PDF Report**

No

**Day(s) Performed**

Monday through Sunday

**Report Available**

Same day/1 to 3 days

**Specimen Retention Time**

7 days

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

**CPT Code Information**

83735

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
MAGU	Magnesium, 24 HR, U	24447-5

Result ID	Test Result Name	Result LOINC® Value
MG24	Magnesium, 24 HR, U	24447-5
TM115	Collection Duration	13362-9
VL111	Urine Volume	3167-4