Test Definition: TTRZ

TTR Gene, Full Gene Analysis, Varies

Overview

Useful For

Establishing a molecular diagnosis for patients with amyloidosis

Identifying variants within *TTR* known to be associated with amyloidosis, allowing for predictive testing of at-risk family members

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULAF	Amniotic Fluid Culture/Genetic Test	Yes	No
_STR1	Comp Analysis using STR (Bill only)	No, (Bill only)	No
_STR2	Add'l comp analysis w/STR (Bill Only)	No, (Bill only)	No
CULFB	Fibroblast Culture for Genetic Test	Yes	No
MATCC	Maternal Cell Contamination, B	Yes	No

Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in one gene associated with amyloidosis: *TTR*. See Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for amyloidosis.

Testing Algorithm

For more information see:
-Amyloidosis (Familial) Test Algorithm
-Amyloidosis: Laboratory Approach to Diagnosis

Special Instructions

- Informed Consent for Genetic Testing
- Amyloidosis: Laboratory Approach to Diagnosis
- Amyloidosis (Familial) Test Algorithm
- Hereditary Peripheral Neuropathy Diagnostic Algorithm
- Molecular Genetics: Neurology Patient Information
- Informed Consent for Genetic Testing (Spanish)

Method Name

Sequence Capture and Targeted Next-Generation Sequencing followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

NY State Available
Yes

Specimen

Specimen Type
Varies

Ordering Guidance
Targeted testing (also called site-specific or known variants testing) is available for variants identified in the *TTR* gene. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

Testing for *TTR* gene as part of a customized panel is available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies.

Specimen Required
Patient Preparation: A previous bone marrow transplant from an allogenic donor will interfere with testing. For instructions for testing patients who have received a bone marrow transplant, call 800-533-1710.

Submit only 1 of the following specimens:

Specimen Type: Whole blood
Container/Tube:
Preferred: Lavender top (EDTA) or yellow top (ACD)
Acceptable: Green top (Sodium heparin):
Specimen Volume: 3 mL
Collection Instructions:
1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot**.
Specimen Stability Information: Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days
Additional Information:
1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA is met, the preferred volume of blood must be submitted. Testing may be canceled if DNA requirements are inadequate.

Specimen Type: Saliva
Patient Preparation: Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.
Supplies:

DNA Saliva Kit High Yield (T1007)
Saliva Swab Collection Kit (T786)
Container/Tube:
Preferred: High-yield DNA saliva kit
Acceptable: Saliva swab
Specimen Volume: 1 Tube if using T1007 or 2 swabs if using T786
Collection Instructions: Collect and send specimen per kit instructions.
Specimen Stability Information: Ambient (preferred) 30 days/Refrigerated 30 days
Additional Information: Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

Specimen Type: Cultured fibroblasts
Source: Skin
Container/Tube: T-25 flask
Specimen Volume: 2 Flasks
Collection Instructions: Submit confluent cultured fibroblast cells from a skin biopsy.
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours
Additional Information:
1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Specimen Type: Tissue biopsy
Supplies: Hank's Solution (T132)
Container/Tube: Sterile container with sterile Hank's balanced salt solution, Ringer's solution, or normal saline
Specimen Volume: 0.5 to 3 cm(3) or larger
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours
Additional Information:
1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.

Specimen Type: Extracted DNA
Container/Tube:
Preferred: Screw Cap Micro Tube, 2 mL with skirted conical base
Acceptable: Matrix tube, 1mL
Collection Instructions:
1. The preferred volume is at least 100 mcL at a concentration of 75 ng/mcL.
2. Include concentration and volume on tube.
Specimen Stability Information: Frozen (preferred) 1year /Ambient/Refrigerated
Additional Information: DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from

a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report

Specimen Type: Blood spot
Supplies: Card-Blood Spot Collection (Filter Paper) (T493)
Container/Tube:
Preferred: Collection card (Whatman Protein Saver 903 Paper)
Acceptable: PerkinElmer 226 filter paper or blood spot collection card
Specimen Volume: 2 to 5 Blood spots
Collection Instructions:
1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see How to Collect a Dried Blood Spot Sample.
2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
3. Do not expose specimen to heat or direct sunlight.
4. Do not stack wet specimens.
5. Keep specimen dry.
Specimen Stability Information: Ambient (preferred)/Refrigerated
Additional Information:
1. Blood spot specimens are acceptable but not recommended. Multiple extractions will be required to obtain sufficient yield for supplemental analysis, and there is significant risk for test failure due to insufficient DNA.
2. Due to lower concentration of DNA yielded from blood spot, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.
3. For collection instructions, see Blood Spot Collection Instructions
4. For collection instructions in Spanish, see Blood Spot Collection Card-Spanish Instructions (T777)
5. For collection instructions in Chinese, see Blood Spot Collection Card-Chinese Instructions (T800)

Prenatal Specimens
Due to its complexity, consultation with the laboratory is required for all prenatal testing; call 800-533-1710 to speak to a genetic counselor.

Specimen Type: Chorionic villi
Container/Tube: 15-mL tube containing 15 mL of transport media
Specimen Volume: 20 mg
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours
Additional Information:
1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Prenatal cultured fibroblasts (eg, products of conception), amniocytes, or other confluent cultured cells. This does not include cultured chorionic villi.
Container/Tube: T-25 flask
Specimen Volume: 2 Flasks
Collection Instructions: Submit confluent cultured cells from another laboratory.
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours
Additional Information:
1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks is required to culture fibroblasts before genetic testing can occur.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Specimen Type: Cultured chorionic villi
Container/Tube: T-25 flasks
Specimen Volume: 2 Full flasks
Collection Instructions: Submit confluent cultured cells from another laboratory.
Specimen Stability Information: Ambient (preferred) <24 hours/Refrigerated <24 hours
Additional Information:
1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing.
3. **All prenatal specimens must be accompanied by a maternal blood specimen;** order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on the maternal specimen.

Forms

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file.
The following documents are available:
-Informed Consent for Genetic Testing (T576)
-Informed Consent for Genetic Testing (Spanish) (T826)
2. Molecular Genetics: Neurology Patient Information
3. If not ordering electronically, complete, print, and send 1 of the following forms with the specimen:
-Cardiovascular Test Request (T724)
-Hematopathology/Cytogenetics Test Request (T726)
-Neurology Specialty Testing Client Test Request (T732)
-Renal Diagnostics Test Request (T830)
-Biochemical Genetics Test Request (T798)

Specimen Minimum Volume

See Specimen Required

Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

Clinical & Interpretive

Clinical Information

The systemic amyloidoses include a number of disorders of varying etiology characterized by extracellular protein deposition. The most common form is an acquired amyloidosis secondary to multiple myeloma or monoclonal gammopathy of unknown significance in which the amyloid is composed of immunoglobulin light chains. In addition to light chain amyloidosis, there are a number of acquired amyloidoses caused by the misfolding and precipitation of a wide variety of proteins. There are also hereditary forms of amyloidosis. Due to the clinical overlap between the acquired and hereditary forms, it is imperative to determine the specific type of amyloidosis in order to provide an accurate prognosis and consider appropriate therapeutic interventions.

The most common hereditary amyloidosis is familial transthyretin amyloidosis, an autosomal dominant disorder caused by variants in the transthyretin (*TTR*) gene. The resulting amino acid substitutions lead to a relatively unstable, amyloidogenic TTR protein. Most individuals begin to exhibit clinical symptoms between the third and seventh decades of life. Typically, TTR-associated amyloidosis is progressive over a course of 5 to 15 years, and the most common cause of death is cardiomyopathy. Affected individuals may present with a variety of symptoms, including peripheral neuropathy, blindness, cardiomyopathy, nephropathy, autonomic nervous dysfunction, or bowel dysfunction.

It is important to note that this assay does not detect variants associated with non-TTR forms of familial amyloidosis.

Reference Values

An interpretive report will be provided.

Interpretation

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.(1) Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

Cautions

Clinical Correlations:
Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:
Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

Deletion/Duplication Analysis:
This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the genes included on the panel may occur. In these instances, genes included in the ordered test are provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. However, copy number variants for genes not listed in the Method Description are typically not reported or interpreted for haploinsufficiency/triplosensitivity. CMACB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the genes included on the panel.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

Reclassification of Variants:
Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare providers to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may

issue an amended report.

Variant Evaluation:
Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.(1) Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

Clinical Reference

1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. Genet Med 2015;17(5):405-424
2. Manganelli F, Fabrizi GM, Luigetti M, Mandich P, Mazzeo A, Pareyson D. Hereditary transthyretin amyloidosis overview. Neurol Sci. 2022:43:595-604. doi:10.1007/s10072-020-04889-2

Performance

Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of *TTR*, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletion-insertions (delins) less than 40 base pairs (bp), above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in *TTR*

There may be regions of *TTR* that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences.(Unpublished Mayo method)

The reference transcript for *TTR* gene is NM_000371.3. Reference transcript numbers may be updated due to transcript re-versioning. Always refer to the final patient report for gene transcript information referenced at the time of testing.

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory

criteria.

PDF Report
No

Day(s) Performed
Varies

Report Available
14 to 21 days

Specimen Retention Time
Whole blood/Saliva: 30 days (if available); Extracted DNA: 3 months; Blood Spot 1 year (if available)

Performing Laboratory Location
Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees
- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification
This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information
81404
88233-Tissue culture, skin, solid tissue biopsy (if appropriate)
88235-Amniotic fluid culture (if appropriate)
81265-Maternal cell contamination (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
TTRZ	TTR Gene, Full Gene Analysis	94225-0

Result ID	Test Result Name	Result LOINC® Value
617754	Test Description	62364-5
617755	Specimen	31208-2
617756	Source	31208-2
617757	Result Summary	50397-9
617758	Result	82939-0

617759	Interpretation	69047-9
618194	Additional Results	82939-0
617760	Resources	99622-3
617761	Additional Information	48767-8
617762	Method	85069-3
617763	Genes Analyzed	48018-6
617764	Disclaimer	62364-5
617765	Released By	18771-6