

Overview

Useful For

Assessing sensitization to various inhalant allergens commonly found in the Great Plains region including Kansas, North Dakota, Nebraska, and South Dakota

Defining the allergen responsible for eliciting signs and symptoms

Identifying allergens:

- Responsible for allergic response and/or anaphylactic episode
- To confirm sensitization prior to beginning immunotherapy
- To investigate the specificity of allergic reactions to insect venom allergens, drugs, or chemical allergens

Profile Information

| Test Id | Reporting Name             | Available Separately | Always Performed |
|---------|----------------------------|----------------------|------------------|
| IGE     | Immunoglobulin E (IgE), S  | Yes                  | Yes              |
| DP      | House Dust Mites/D.P., IgE | Yes                  | Yes              |
| DF      | House Dust Mites/D.F., IgE | Yes                  | Yes              |
| CAT     | Cat Epithelium, IgE        | Yes                  | Yes              |
| DOGD    | Dog Dander, IgE            | Yes                  | Yes              |
| BERG    | Bermuda Grass, IgE         | Yes                  | Yes              |
| TIMG    | Timothy Grass, IgE         | Yes                  | Yes              |
| COCR    | Cockroach, IgE             | Yes                  | Yes              |
| PENL    | Penicillium, IgE           | Yes                  | Yes              |
| CLAD    | Cladosporium, IgE          | Yes                  | Yes              |
| ASP     | Aspergillus Fumigatus, IgE | Yes                  | Yes              |
| ALTN    | Alternaria Tenuis, IgE     | Yes                  | Yes              |
| BXMPL   | Box Eld/Maple, S, IgE      | Yes                  | Yes              |
| CED     | Mountain Cedar, IgE        | Yes                  | Yes              |
| OAK     | Oak, IgE                   | Yes                  | Yes              |
| ELM     | Elm, IgE                   | Yes                  | Yes              |
| CTWD    | Cottonwood, IgE            | Yes                  | Yes              |
| ASHW    | White Ash, IgE             | Yes                  | Yes              |
| MULB    | Mulberry, IgE              | Yes                  | Yes              |
| SRW     | Short Ragweed, IgE         | Yes                  | Yes              |
| RUSS    | Russian Thistle, IgE       | Yes                  | Yes              |
| SORR    | Red Sorrel, IgE            | Yes                  | Yes              |
| NETT    | Nettle, IgE                | Yes                  | Yes              |

Special Instructions

- [Allergens - Immunoglobulin E \(IgE\) Antibodies](#)

Method Name

Fluorescence Enzyme Immunoassay (FEIA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Ordering Guidance

For a listing of allergens available for testing, see [Allergens - Immunoglobulin E \(IgE\) Antibodies](#)

Specimen Required

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1.6 mL

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms

[If not ordering electronically, complete, print, and send an Allergen Test Request](#) (T236) with the specimen.

Specimen Minimum Volume

1.4 mL

Reject Due To

|                 |    |
|-----------------|----|
| Gross hemolysis | OK |
| Gross lipemia   | OK |

Specimen Stability Information

| Specimen Type | Temperature              | Time    | Special Container |
|---------------|--------------------------|---------|-------------------|
| Serum         | Refrigerated (preferred) | 14 days |                   |
|               | Frozen                   | 90 days |                   |

Clinical & Interpretive

Clinical Information

Immunoglobulin E (IgE) is one of the 5 classes of immunoglobulins and is defined by the presence of the epsilon heavy chain. It is the most recently described immunoglobulin, having first been identified in 1966. IgE exists as a monomer and is present in the circulation at very low concentrations, approximately 300-fold lower than that of IgG. The physiologic role of IgE is not well characterized, although it is thought to be involved in defense against parasites, specifically helminths.

The function of IgE is distinct from other immunoglobulins in that it induces activation of mast cells and basophils through the cell-surface receptor Fc epsilon RI. Fc epsilon RI is a high-affinity receptor specific for IgE present at a high density on tissue-resident mast cells and basophils. Because of this high-affinity interaction, almost all IgE produced by B cells is bound to mast cells or basophils, which explains the low concentration present in circulation. Cross-linking of the Fc epsilon RI-bound IgE leads to cellular activation, resulting in immediate release of preformed granular components (histamine and tryptase) and subsequent production of lipid mediators (prostaglandins and leukotrienes) and cytokines (interleukin-4 and interleukin-5).

Elevated concentrations of IgE may occur in the context of allergic disease. However, increases in the amount of circulating IgE can also be found in various other diseases, including primary immunodeficiencies, infections, inflammatory diseases, and malignancies. Total IgE measurements have limited utility for diagnostic evaluation of patients with suspected allergic disease. In this scenario, testing for the presence of allergen-specific IgE may provide more information.

Clinical manifestations of allergic disease result from activation of mast cells and basophils, which occurs when Fc epsilon RI-bound IgE antibodies interact with allergen.

In vitro serum testing for specific IgE antibodies may provide an indication of the immune response to an allergen that may be associated with allergic disease.

The allergens chosen for testing often depend upon the age of the patient, history of allergen exposure, season of the year, and clinical manifestations. Sensitization to inhalant allergens (dust mites, mold, and pollen inhalants) primarily occurs in older children, adolescents, and adults and usually manifests as respiratory disease (rhinitis and asthma).

Reference Values

Specific IgE:

| Class | IgE kU/L  | Interpretation       |
|-------|-----------|----------------------|
| 0     | <0.10     | Negative             |
| 0/1   | 0.10-0.34 | Borderline/equivocal |
| 1     | 0.35-0.69 | Equivocal            |
| 2     | 0.70-3.49 | Positive             |
| 3     | 3.50-17.4 | Positive             |
| 4     | 17.5-49.9 | Strongly positive    |
| 5     | 50.0-99.9 | Strongly positive    |

|   |           |                   |
|---|-----------|-------------------|
| 6 | > or =100 | Strongly positive |
|---|-----------|-------------------|

Reference values apply to all ages.

Total IgE:

| Age                | Reference interval (in kU/L) |
|--------------------|------------------------------|
| 0-5 months         | < or =13                     |
| 6-11 months        | < or =34                     |
| 1 and 2 years      | < or =97                     |
| 3 years            | < or =199                    |
| 4-6 years          | < or =307                    |
| 7 and 8 years      | < or =403                    |
| 9-12 years         | < or =696                    |
| 13-15 years        | < or =629                    |
| 16 and 17 years    | < or =537                    |
| 18 years and older | < or =214                    |

Interpretation

Elevated concentrations of total IgE may be found in a variety of clinical diseases, including allergic disease, certain primary immunodeficiencies, infections, inflammatory diseases, and malignancies.

Detection of allergen-specific IgE antibodies in serum (class 1 or greater) indicates an increased likelihood of allergic disease as opposed to other etiologies and defines the allergens that may be responsible for eliciting signs and symptoms.

Cautions

An elevated concentration of total IgE is not diagnostic for allergic disease, and it must be interpreted in the clinical context of the patient, including age, sex, travel history, potential allergen exposure, and family history.

A normal concentration of total IgE does not eliminate the possibility of allergic disease. In patients with a high index of suspicion for allergic disease, testing for allergen-specific IgE may be warranted.

Testing for allergen-specific IgE antibodies is not useful in patients previously treated with immunotherapy to determine if residual clinical sensitivity exists or in patients in whom the medical management does not depend upon identification of allergen specificity.

Some individuals with clinically insignificant sensitivity to allergens may have measurable levels of IgE antibodies in serum, and test results must be interpreted in the clinical context.

False-positive results for IgE antibodies may occur in patients with markedly elevated serum IgE (>2500 kU/L) due to nonspecific binding to allergen solid phases.

Clinical Reference

1. Homburger HA, Hamilton RG: Allergic diseases. In: McPherson RA, Pincus MR, eds. Henry's Clinical Diagnosis and Management by Laboratory Methods. 23rd ed. Elsevier; 2017:1057-1070

2. Bernstein IL, Li JT, Bernstein DI, et al: Allergy diagnostic testing: an updated practice parameter. Annals Allergy Asthma and Immunol 2008; 100:S1-S148

Performance

Method Description

Specific IgE:  
Specific IgE from the patient's serum reacts with the allergen of interest, which is covalently coupled to an ImmunoCAP. After washing away nonspecific IgE, enzyme-labeled anti-IgE antibody is added to form a complex. After incubation, unbound anti-IgE is washed away, and the bound complex incubated with a developing agent. After stopping the reaction, the fluorescence of the eluate is measured. Fluorescence is proportional to the amount of specific IgE present in the patient's sample (ie, the higher the fluorescence value, the more IgE antibody is present).(Package insert: ImmunoCAP System Specific IgE FEIA. Phadia; Rev 06/2020)

Total IgE:  
Anti-IgE, covalently coupled to ImmunoCAP, reacts with the IgE in a serum specimen. After washing, enzyme-labeled anti-IgE antibodies are added to form a complex. After incubation, unbound enzyme-labeled anti-IgE is washed away, and the bound complex incubated with a developing agent. After stopping the reaction, fluorescence of the eluate in the well is measured. The fluorescence is directly proportional to the concentration of IgE in the test specimen.(Package insert: Phadia CAP System IgE FEIA. Phadia; Rev 10/2019)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

Same day/1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.

- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

82785-IgE  
86003 x 22-Each individual allergen

LOINC® Information

| Test ID | Test Order Name                   | Order LOINC® Value |
|---------|-----------------------------------|--------------------|
| RPR9    | Resp Profile, Reg 9, Great Plains | 94593-1            |

| Result ID | Test Result Name           | Result LOINC® Value |
|-----------|----------------------------|---------------------|
| ALTN      | Alternaria Tenuis, IgE     | 6020-2              |
| ASHW      | White Ash, IgE             | 6278-6              |
| ASP       | Aspergillus Fumigatus, IgE | 6025-1              |
| BERG      | Bermuda Grass, IgE         | 6041-8              |
| BXMPL     | Box Eld/Maple, S, IgE      | 7155-5              |
| CAT       | Cat Epithelium, IgE        | 6833-8              |
| CED       | Mountain Cedar, IgE        | 6178-8              |
| CLAD      | Cladosporium, IgE          | 53760-5             |
| COCR      | Cockroach, IgE             | 6078-0              |
| CTWD      | Cottonwood, IgE            | 6090-5              |
| DF        | House Dust Mites/D.F., IgE | 6095-4              |
| DOGD      | Dog Dander, IgE            | 6098-8              |
| DP        | House Dust Mites/D.P., IgE | 6096-2              |
| ELM       | Elm, IgE                   | 6109-3              |
| IGE       | Immunoglobulin E (IgE), S  | 19113-0             |
| MULB      | Mulberry, IgE              | 6281-0              |
| NETT      | Nettle, IgE                | 6186-1              |
| OAK       | Oak, IgE                   | 6189-5              |
| PENL      | Penicillium, IgE           | 6212-5              |
| RUSS      | Russian Thistle, IgE       | 6234-9              |
| SORR      | Red Sorrel, IgE            | 6244-8              |
| SRW       | Short Ragweed, IgE         | 6085-5              |
| TIMG      | Timothy Grass, IgE         | 6265-3              |