

Overview

Useful For

Diagnosis of acute or recent hepatitis A infection

Special Instructions

- [Viral Hepatitis Serologic Profiles](#)

Method Name

Electrochemiluminescence Immunoassay (ECLIA)

NY State Available

No

Specimen

Specimen Type

Serum SST

Necessary Information

Date of collection is required.

Specimen Required

Patient Preparation: For 24 hours before specimen collection, patient **should not** take multivitamins or dietary supplements (eg, hair, skin, and nail supplements) containing biotin (vitamin B7).

Collection Container/Tube: Serum gel (red-top tubes are **not acceptable**)

Submission Container/Tube: Plastic vial

Specimen Volume: 0.6 mL

Collection Instructions:

1. Centrifuge blood collection tube per manufacturer's instructions (eg, centrifuge and aliquot within 2 hours of collection for BD Vacutainer tubes).
2. Aliquot serum into plastic vial.

Forms

If not ordering electronically, complete, print, and send 1 of the following:

-[Gastroenterology and Hepatology Test Request](#) (T728)

-[Infectious Disease Serology Test Request](#) (T916)

Specimen Minimum Volume

0.6 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum SST	Frozen (preferred)	90 days	
	Refrigerated	6 days	
	Ambient	72 hours	

Clinical & Interpretive**Clinical Information**

Hepatitis A virus (HAV) is endemic throughout the world, occurring most commonly in areas of poor hygiene and low socioeconomic conditions. The virus is transmitted primarily by the fecal-oral route and spread by close person-to-person contact and by food and waterborne epidemics. Outbreaks frequently occur in overcrowded situations and high-density institutions and centers, such as prisons and healthcare or daycare centers. Viral spread by parenteral routes (eg, exposure to blood) is possible, but rare, because infected individuals are viremic for a short period of time (usually <3 weeks). There is little or no evidence of transplacental transmission from mother to fetus or transmission to newborn during delivery.

Serological diagnosis of acute viral hepatitis A depends on the detection of specific anti-HAV IgM. Its presence in the patient's serum indicates a recent exposure to HAV. HAV-specific IgM antibody level becomes detectable in the blood by 4 weeks after infection, persisting at elevated levels for about 2 months before declining to undetectable levels by 6 months. They rarely persist beyond 12 months after infection.

Reference Values

Negative

See [Viral Hepatitis Serologic Profiles](#).

Interpretation

This assay detects the presence of hepatitis A virus (HAV)-specific IgM antibody in serum.

Negative results indicate either inadequate or delayed anti-HAV IgM response after known exposure to HAV or absence of acute or recent hepatitis A.

Equivocal results may be seen in early acute hepatitis A associated with rising anti-HAV IgM levels or recent hepatitis A infection associated with declining anti-HAV IgM levels. Retesting for both anti-HAV IgM (HAIGM / Hepatitis A Virus IgM Antibody, Serum) and anti-HAV Total (HAVTA / Hepatitis A Virus Total Antibodies, Serum) in 2 to 4 weeks is recommended to determine the definitive HAV infection status.

Positive results indicate acute or recent (<6 months) hepatitis A infection. As required by laws in almost all states, positive anti-HAV IgM test results must be urgently reported to state health departments for epidemiologic investigations of possible outbreak transmission.

Cautions

Testing too early (<2 weeks) after exposure to hepatitis A virus (HAV) may yield negative anti-HAV IgM results.

False-positive results may be due to presence of cross-reactive antibodies from other viral infection or underlying illnesses (such as non-Hodgkin lymphoma). Positive results should be correlated with patient's clinical history and epidemiologic exposure.

The presence of heterophilic antibodies and human antimouse antibodies (in patients who have received preparations of mouse monoclonal antibodies for diagnosis or therapy) in serum may interfere with the assay and cause erroneous results (false-positive or false-negative).

Consumption of high-dose biotin supplement within 12 hours of blood collection for this test can cause false-negative test results. Individuals should cease taking these biotin-containing dietary supplements for minimum 12 hours before blood collection for this test.

Performance characteristics have not been established for the following specimen characteristics:

- Grossly icteric (total bilirubin level of >50 mg/dL)
- Grossly hemolyzed (hemoglobin level of >1000 mg/dL)
- Grossly lipemic (intralipid >2000 mg/dL)
- Containing particulate matter
- Heat-inactivated
- Cadaveric specimens

Clinical Reference

1. de Paula VS. Laboratory diagnosis of hepatitis A. *Future Virology*. 2012;7(5):461-472
2. Nelson NP, Weng MK, Hofmeister MG, et al. Prevention of hepatitis A infection in the United States: Recommendations of the Advisory Committee on Immunization Practices, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(5):1-38. Erratum in *MMWR Morb Mortal Wkly Rep*. 2021;70(8):294
3. Webb GW, Kelly S, Dalton HR. Hepatitis A and hepatitis E: clinical and epidemiological features, diagnosis, treatment, and prevention. *Clin Microbiol Newslett*. 2020;42(21):171-179

Performance**Method Description**

The Elecsys Anti-HAV (hepatitis A virus) IgM assay is performed using an electrochemiluminescence immunoassay on the automated cobas e 801 immunochemistry analyzer. Hepatitis A virus-specific IgM antibody (anti-HAV IgM) in the patient's serum sample is pretreated with anti-Fdy reagent to block specific IgG in the presence of monoclonal anti-HAV antibodies labeled with ruthenium complex. After addition of biotinylated monoclonal h-IgM-specific antibodies, HAV antigen, and streptavidin-coated microparticles, patient's anti-HAV IgM form a sandwich complex with the HAV antigen

and the ruthenium-labeled anti-HAV antibody which becomes bound to the solid phase via interaction of biotin and streptavidin. The reaction mixture is then aspirated into the measuring cell where the microparticles are magnetically captured onto the surface of the electrode, and unbound substances are washed away. Voltage is applied to the electrode then induces chemiluminescent emissions that are measured by a photomultiplier. Test result for each patient's sample is determined automatically by the assay-specific software program by comparing the electrochemiluminescence signal generated from the patient's sample to the cutoff index value set from reagent lot-specific assay calibrations. (Package insert: Elecsys Anti-HAV IgM. Roche Diagnostics; v5.0, 11/2022)

PDF Report

No

Day(s) Performed

Monday through Friday, Sunday

Report Available

1 to 2 days

Specimen Retention Time

14 days

Performing Laboratory Location

Jacksonville

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

86709

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
HAIGM	Hepatitis A IgM Ab, S	13950-1

Result ID	Test Result Name	Result LOINC® Value
HAIGM	Hepatitis A IgM Ab, S	13950-1