



Test Definition: RTSCJ

Reptilase Time, Plasma

Overview

Useful For

Evaluation of a prolonged thrombin time (TT)

Confirm or exclude the presence of heparin in the specimen or specimen type

Evaluating hypofibrinogenemia or dysfibrinogenemia in conjunction with the TT and fibrinogen assay

Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)

Method Name

Optical Clot-Based

NY State Available

No

Specimen

Specimen Type

Plasma Na Cit

Specimen Required

Specimen Type: Platelet-poor plasma

Collection Container/Tube: Light-blue top (3.2% sodium citrate)

Submission Container/Tube: Plastic vial

Specimen Volume: 1 mL

Collection Instructions:

1. For complete instruction, see [Coagulation Guidelines for Specimen Handling and Processing](#).
2. Centrifuge, transfer all plasma into a plastic vial, and centrifuge plasma again.
3. Aliquot plasma into a plastic vial, leaving 0.25 mL in the bottom of centrifuged vial.
4. Freeze plasma immediately (no longer than 4 hours after collection) at -20 degrees C or, ideally below -40 degrees C.

Additional Information:

1. Double-centrifuged specimen is critical for accurate results as platelet contamination may cause spurious results.
2. Each coagulation assay requested should have its own vial.

Specimen Minimum Volume

0.5 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Clinical & Interpretive

Clinical Information

Prolonged clotting times may be associated with a wide variety of coagulation abnormalities including:

- Deficiency or functional abnormality (congenital or acquired) of any of the coagulation proteins
- Deficiency or functional abnormality of platelets
- Specific factor inhibitors
- Acute disseminated intravascular coagulation
- Exogenous anticoagulants (eg, heparin, warfarin)

The prothrombin time (PT) and activated partial thromboplastin time (aPTT) are first-order tests for coagulation abnormalities and are prolonged in many bleeding disorders. A battery of coagulation tests is often required to determine the cause of prolonged clotting times. The thrombin time (TT) test is used to identify the cause of prolonged aPTT or dilute Russell's viper venom time (DRVVT). Reptilase time (RT) test is used to evaluate a prolonged TT.

Reptilase is a thrombin-like enzyme isolated from the venom of *Bothrops atrox*. Thrombin splits small fibrinopeptides A and B from fibrinogen molecules, producing fibrin monomer, which polymerizes to form a clot. Reptilase, however, splits off fibrinopeptide A but not B, which results in fibrin polymerization. In contrast to thrombin and the TT test which are inhibited by heparin, the RT is normal in the presence of heparin. Similar to the TT test, the RT is prolonged in the presence of hypofibrinogenemia and dysfibrinogenemia.

Reference Values

14.0-23.9 seconds

Interpretation

As seen in the following table, reptilase time can help distinguish among the various causes of a prolonged thrombin time (TT).

Table.

Thrombin time	Reptilase time	Causes	Remarks
Prolonged	Prolonged	Hypo- or afibrinogenemia	Ascertain by determination of fibrinogen
Prolonged	Prolonged	Dysfibrinogenemia	Ascertain by specific assay

Prolonged	Normal	Heparin or inhibitor of thrombin	Differentiate by human TT and/or heparin assays
Prolonged	Prolonged	Fibrin(ogen) split products (FSP)	Ascertain by FSP or D-dimer assay

Cautions

The reptilase time test has limited diagnostic value when ordered as a stand-alone test.

Clinical Reference

Favaloro EJ, Lippi G, eds. Hemostasis and Thrombosis: Methods and Protocols. Humana Press; 2017

Performance

Method Description

Reptilase time is used in addition to thrombin time to exclude sample heparin contamination in cases of prolonged thrombin time. Reptilase is a serine protease from the venom of the Bothrops atrox snake (batroxobin), mimicking thrombin and cleaving fibrinogen's fibrinopeptide A. In contrast, thrombin cleaves both fibrinopeptides A and B. Adding reptilase to platelet-poor plasma activates thrombus formation, which is detected using optical or electro-mechanical methods. (Danilatou V. Laboratory Evaluation of Coagulopathies. In Zubair M, ed. StatPearls Online. Updated July 27, 2024. Accessed March 26, 2025. Available at www.statpearls.com/point-of-care/145925)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 4 days

Specimen Retention Time

14 days

Performing Laboratory Location

Mayo Clinic Jacksonville Clinical Lab

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

CPT Code Information

85635

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
RTSCJ	Reptilase Time, P	6683-7

Result ID	Test Result Name	Result LOINC® Value
RTSCJ	Reptilase Time, P	6683-7