

Clonazepam and 7-Aminoclonazepam, Serum

## **Overview**

#### **Useful For**

Assessing patient compliance

Monitoring for appropriate therapeutic level

Assessing clonazepam toxicity

#### **Method Name**

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

#### **NY State Available**

Yes

## **Specimen**

## **Specimen Type**

Serum Red

#### **Specimen Required**

Collection Container/Tube: Red top Submission Container/Tube: Plastic vial

**Specimen Volume:** 1.2 mL **Collection Instructions:** 

- 1. Draw blood immediately before next scheduled dose (minimum 12 hours after last dose).
- 2. Within 2 hours of collection, the specimen must be centrifuged and the serum aliquoted into a plastic vial.

#### **Forms**

If not ordering electronically, complete, print, and send a <u>Therapeutics Test Request</u> (T831) with the specimen.

## **Specimen Minimum Volume**

0.6 mL

#### **Reject Due To**

| Gross         | ОК     |
|---------------|--------|
| hemolysis     |        |
| Gross lipemia | ОК     |
| Gross icterus | Reject |



## Clonazepam and 7-Aminoclonazepam, Serum

## **Specimen Stability Information**

| Specimen Type | Temperature              | Time     | Special Container |
|---------------|--------------------------|----------|-------------------|
| Serum Red     | Refrigerated (preferred) | 14 days  |                   |
|               | Frozen                   | 28 days  |                   |
|               | Ambient                  | 72 hours |                   |

## Clinical & Interpretive

#### **Clinical Information**

Clonazepam (5 [2-chlorophenyl]-2, 3-dihydro-7-nitro-1, 4-benzodiazepin-2-one), a benzodiazepine is useful alone or as an adjunct in the treatment of certain seizures. In addition, it may be useful in patients with panic disorder and restless legs syndrome.

Clonazepam has no definite antiseizure and antipanic mechanism of action, although it is believed to be related to its capacity to enhance gamma-aminobutyric acid (GABA) activity, which is the major inhibitory neurotransmitter in the central nervous system. It is able to suppress the spike and wave discharges in absence seizures and decreases the frequency, duration, amplitude, and spread of discharge in minor motor seizures.

Clonazepam is highly protein bound (approximately 85%). It is extensively metabolized by hepatic cytochrome P450, family 3, subfamily A (CYP3A) to inactive metabolites and has a half-life of 30 to 40 hours.

## **Reference Values**

Clonazepam

Anticonvulsant: 20-70 ng/mL

Anxiolytic: 4-80 ng/mL

Some individuals may show therapeutic response outside of these ranges or may display toxicity within the therapeutic range, thus interpretation should include clinical evaluation.

**Note:** Therapeutic ranges are for specimens collected at trough (ie, immediately before next scheduled dose). Levels may be elevated in non-trough specimens.

#### Interpretation

The therapeutic range varies depending on the indication.

Some individuals may respond well outside of these ranges or may display toxicity within the therapeutic range, and thus, interpretation should include clinical evaluation.

The possibility of toxicity is increased when levels exceed 100 ng/mL.

#### **Cautions**

Specimens that are obtained from gel tubes must be removed from the gel within 2 hours.

## **Clinical Reference**

Hiemke C, Baumann P, Bergemann N, et al: AGNP Consensus Guidelines for Therapeutic Drug Monitoring in Psychiatry: Update 2011. Pharmacopsychiatry. 2011;44(6):195-235



Clonazepam and 7-Aminoclonazepam, Serum

#### **Performance**

## **Method Description**

Clonazepam and 7-aminoclonazepam are extracted from serum using a liquid-liquid extraction technique. The organic layer is removed, dried, and reconstituted. Analysis is by liquid chromatography-tandem mass spectrometry. (Unpublished Mayo method)

#### **PDF Report**

No

## Day(s) Performed

Monday, Wednesday

#### Report Available

3 to 7 days

## **Specimen Retention Time**

14 days

#### **Performing Laboratory Location**

Rochester

## **Fees & Codes**

#### **Fees**

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

## **Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

## **CPT Code Information**

80346

G0480 (if appropriate)

#### **LOINC®** Information

| Test ID | Test Order Name      | Order LOINC® Value |
|---------|----------------------|--------------------|
| CZPS    | Clonazepam and       | In Process         |
|         | 7-Aminoclonazepam, S |                    |



Clonazepam and 7-Aminoclonazepam, Serum

| Result ID | Test Result Name  | Result LOINC® Value |
|-----------|-------------------|---------------------|
| 65044     | Clonazepam        | 3494-2              |
| 41782     | 7-Aminoclonazepam | 28059-4             |