

Test Definition: CTXP

Cerebrotendinous Xanthomatosis, Plasma

Overview

Useful For

Evaluating patients with a clinical suspicion of cerebrotendinous xanthomatosis (CTX) using plasma specimens

Monitoring of individuals with CTX on chenodeoxycholic acid (CDCA) therapy

This test is not useful for the identification of carriers

This test is not useful for the evaluation of bile acid malabsorption

Special Instructions

Biochemical Genetics Patient Information

Method Name Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)

NY State Available Yes

Specimen

Specimen Type Plasma

Ordering Guidance

For assessment of bile acid malabsorption in patients with irritable bowel syndrome-diarrhea, order 7AC4 / 7AC4, Bile Acid Synthesis, Serum.

This test is also available as a part of a panel; see HSMP / Hepatosplenomegaly Panel, Plasma. If this test (CTXP) is ordered with either GPSYP / Glucopsychosine, Plasma or OXNP / Oxysterols, Plasma, the individual tests will be canceled and HSMP ordered.

Specimen Required

Collection Container/Tube: Preferred: Lavender top (EDTA) Acceptable: Green top (sodium heparin, lithium heparin), yellow top (ACD B) Submission Container/Tube: Plastic vial Specimen Volume: 0.3 mL Collection Instructions: 1. Centrifuge at 4 degrees C if possible



Test Definition: CTXP

Cerebrotendinous Xanthomatosis, Plasma

2. Aliquot plasma into a plastic vial. **Do not disturb or transfer the buffy coat layer**.

3. Send frozen.

Forms

1. Biochemical Genetics Patient Information (T602)

2. If not ordering electronically, complete, print, and send a <u>Biochemical Genetics Test Request</u> (T798) with the specimen.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross	ОК
hemolysis	
Gross lipemia	ОК
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma	Frozen	65 days	

Clinical & Interpretive

Clinical Information

Cerebrotendinous xanthomatosis (CTX) is an autosomal recessive disorder of bile acid synthesis resulting from the deficiency of the mitochondrial enzyme, sterol 27-hydrolase. Sterol 27-hydrolase facilitates the first step of sterol degradation in the formation of bile acids; consequently, patients with CTX will experience increased storage of the sterol, cholestenol, and ketosterol bile acid precursors (7-alpha-hydroxy-4-cholesten-3-one [7a-C4] and 7-alpha,12-alpha–dihydroxycholest-4-en-3-one [7a12aC4]) in multiple tissues throughout the body with a resulting deficiency of the bile acid, chenodeoxycholic acid (CDCA). CTX is caused by variants in the *CYP27A1* gene.

Patients with CTX can present with a constellation of findings, including infantile onset diarrhea, childhood onset cataracts, development of tendon/cerebral xanthomas in adolescence and early adulthood, early onset osteoporosis, as well as a broad array of neuropsychological manifestations, such as intellectual disability, dementia, psychiatric symptoms, ataxia, pyramidal signs, dystonia, and muscle weakness. Patients may occasionally present with cholestatic liver disease, which may present as jaundice, poor growth, and hepatosplenomegaly. Intrafamilial variability exists and some heterozygous carriers may experience a higher incidence of cardiac disorders or gallstones. Treatment with CDCA normalizes bile acid synthesis and suppresses cholestenol biosynthesis, with improvement of clinical symptoms and arrest of disease progression. However, more recently, cholic acid has been proposed as an alternative treatment for adults with CTX. Supplementation with beta-hydroxy beta-methylglutaryl-CoA (HMG-CoA) inhibitors and coenzyme Q10 has also been proposed. The availability of effective therapy makes early diagnosis and treatment of patients with CTX essential.



Cerebrotendinous Xanthomatosis, Plasma

The estimated incidence of CTX is 1 in 50,000 individuals of Northern European ancestry and as high as 1 in 440 in the Druze population of Israel.

The diagnostic evaluation of patients with suspected CTX may reveal abnormalities on brain magnetic resonance imaging (eg, cerebellar atrophy, decrease in volume of grey and white matter, and abnormal white matter signal) in addition to the biochemical and clinical abnormalities. The biochemical diagnosis of CTX can be confirmed by molecular genetic analysis of the *CYP27A1* gene (included in CHLGP / Cholestasis Gene Panel, Varies; or order CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies and indicate the gene to be assessed).

Reference Values

7-ALPHA-HYDROXY-4-CHOLESTEN-3-ONE (7a-C4) Cutoff: < or =0.300 nmol/mL

7-ALPHA,12 ALPHA–DIHYDROXYCHOLEST-4-en-3-ONE (7a12aC4) Cutoff: < or =0.100 nmol/mL

Interpretation

An elevation of 7-alpha-hydroxy-4-cholesten-3-one (7a-C4) and 7-alpha,12-alpha-dihydroxycholest-4-en-3-one (7a12aC4) is strongly suggestive of cerebrotendinous xanthomatosis.

Cautions

Patients with bile acid malabsorption or ileal resection may have elevations of 7-alpha-hydroxy-4-cholesten-3-one (7aC4).

Clinical Reference

1. Mignarri A, Magni A, Del Puppo M, et al: Evaluation of cholesterol metabolism in cerebrotendinous xanthomatosis. J Inherit Metab Dis. 2016 Jan;39(1):75-83

2. Nie S, Chen G, Cao X, Zhang Y: Cerebrotendinous xanthomatosis: a comprehensive review of pathogenesis, clinical manifestations, diagnosis, and management. Orphanet J Rare Dis. 2014 Nov 26;9:179

3. DeBarber AE, Luo J, Star-Weinstock M, et al: A blood test for cerebrotendinous xanthomatosis with potential for disease detection in newborns. J Lipid Res. 2014 Jan;55(1):146-154

 Federico A, Gallus GN: Cerebrotendinous xanthomatosis. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. GeneReviews [Internet]. University of Washington, Seattle; 2003. Updated March 17, 2022. Accessed December 28, 2022. Available at www.ncbi.nlm.nih.gov/books/NBK1409/

5. Lutjohann D, Stellaard F, Bjorkhem I: Levels of 7alpha-hydroxycholesterol and/or 7alpha-hydroxy-4-cholest-3-one are the optimal biochemical markers for the evaluation of treatment of cerebrotendinous xanthomatosis. J Neurol. 2020 Feb;267(2):572-573. doi: 10.1007/s00415-019-09650-0

6. Mandia D, Chaussenot A, Besson G, et al: Cholic acid as a treatment for cerebrotendinous xanthomatosis in adults. J Neurol. 2019 Aug;266(8):2043-2050. doi: 10.1007/s00415-019-09377-y

Performance

Method Description



Test Definition: CTXP

Cerebrotendinous Xanthomatosis, Plasma

Internal standard is added to an aliquot of plasma which is then subjected to protein precipitation. Following centrifugation, the supernatant is subjected to liquid chromatography tandem mass spectrometry (LC-MS/MS) analysis. The MS/MS is operated in the multiple reaction monitoring positive mode to follow the precursor to product species transitions for each analyte and internal standard. The ratio of the extracted peak areas to internal standard determined by the LC-MS/MS is used to calculate the concentration of in the sample.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed Tuesday, Thursday

Report Available 3 to 7 days

Specimen Retention Time 2 months

Performing Laboratory Location Rochester

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82542

LOINC[®] Information

Test ID	Test Order Name	Order LOINC [®] Value	
СТХР	Cerebrotendinous Xanthomatosis, P	92746-7	
Result ID	Test Result Name	Result LOINC [®] Value	
BA4379	Interpretation (CTXP)	59462-2	
BA4376	7a-hydroxy-4-cholesten-3-one	92761-6	
BA4377	7a,12a-dihydroxycholest-4-en-3-one	92758-2	
BA4378	Reviewed By	18771-6	