

Overview

Useful For

As a component to the initial evaluation of a patient presenting with hepatosplenomegaly, using plasma specimens

This test is **not useful for** the identification of carriers.

This test **should not be used** as a monitoring tool for patients with confirmed diagnoses.

Highlights

This is a screening test for a select number of lysosomal and lipid storage disorders, including cerebrotendinous xanthomatosis, Gaucher disease, and Niemann-Pick disease types A, B (also known as acid sphingomyelinase deficiency), and C.

The above conditions may all have hepatosplenomegaly as a presenting sign, making this test a helpful component of a patient's initial evaluation.

Method Name

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

NY State Available

Yes

Specimen

Specimen Type

Plasma

Ordering Guidance

This test **should not be used** for monitoring of patients with confirmed diagnoses. If testing requested is for monitoring purposes, see:

CTXP / Cerebrotendinous Xanthomatosis, Plasma

GPSYP / Glucopsychosine, Plasma

OXNP / Oxysterols, Plasma

Specimen Required

Collection Container/Tube:

Preferred: Lavender top (EDTA)

Acceptable: Green top (sodium heparin, lithium heparin), yellow top (ACD B)

Submission Container/Tube: Plastic vial

Specimen Volume: 0.3 mL

Collection Instructions:

1. Centrifuge at 4 degrees C, if possible
2. Aliquot plasma into plastic vial. **Do not disturb or transfer the buffy coat layer.**
3. Send frozen

Forms

[If not ordering electronically, complete, print, and send a Biochemical Genetics Test Request](#) (T798) with the specimen.

Specimen Minimum Volume

0.25 mL

Reject Due To

Gross hemolysis	OK
Gross lipemia	OK
Gross icterus	OK

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma	Frozen	65 days	

Clinical & Interpretive

Clinical Information

Hepatosplenomegaly is a presenting or accompanying feature for many different inborn errors of metabolism. It typically is a consequence of chronic hepatic dysfunction or abnormal storage of lipids, sugars, or other improperly metabolized analytes due to a particular enzymatic deficiency. The diagnosis can occasionally be narrowed down by consideration of clinical symptoms; however, clinical diagnosis can be difficult due to similarity of clinical features across disorders as well as phenotypic variability. Therefore, screening tests can play an important role in the workup of a patient presenting with hepatosplenomegaly who may have a lysosomal or lipid storage disorder.

The conditions detected in this assay are cerebrotendinous xanthomatosis, Gaucher disease, and Niemann-Pick disease types A, B (also known as acid sphingomyelinase deficiency), and C.

Patients with abnormal results should have follow-up enzymatic or molecular testing for confirmation of diagnosis.

Table. Conditions Identifiable by Method

Disorder	Onset	Analyte detected	Gene	Incidence
Cerebrotendinous xanthomatosis (CTX)	Infancy-adulthood	7-Alpha-hydroxy-4-cholesten-3-one (7a-C4) 7-Alpha,12-alpha-dihydroxycholest-4-en-3-one	CYP27A1	1 in 50,000 As high as 1 in 400 in Druze population.

		(7a12aC4)		
	Phenotype: Early onset diarrhea, cataracts, tendon/cerebral xanthomas, osteoporosis, neuropsychological manifestations, liver disease/hepatosplenomegaly.			
Gaucher disease	Type I: childhood/adult	Glucopsychosine (GPSY)	GBA	Type I: 1 in 30,000 to 1 in 100,000
	Types II/III: neonatal-early childhood			Types II/III: 1 in 100,000
	Phenotype: All types exhibit hepatosplenomegaly and hematological abnormalities. Type I: Organomegaly, thrombocytopenia, and bone pain. Absence of neurologic symptoms. Types II/III: Primary neurologic disease, developmental delay/regression, hepatosplenomegaly, lung disease. Patients with type II typically die by 2 to 4 years of age. Patients with type III may have a less progressive phenotype and may survive into adulthood.			
Niemann-Pick type A/B (NPA/NPB)	NPA: neonatal	Lyso-sphingomyelin (LSM)	SMPD1	Combined incidence
	NPB: birth-adulthood	LSM 509		1 in 250,000
	Phenotype: NPA: Feeding difficulties, jaundice, hepatosplenomegaly, neurologic deterioration, lung disease, hearing and vision impairment, cherry red macula, death usually by 3 years of age. NPB: Mainly limited to visceral symptoms; hepatosplenomegaly, stable liver dysfunction, pulmonary compromise, osteopenia.			
Niemann-Pick type C (NPC)	Variable (perinatal-adulthood)	Cholestane-3 beta, 5-alpha, 6-beta-triol (COT)	NPC1 or NPC2	1 in 120,000 to 1 in 150,000
	Phenotype: Variable clinical presentation; ataxia, vertical supranuclear gaze palsy, dystonia, progressive speech deterioration, seizures, +/- hepatosplenomegaly.			

Patients with testing indicative of one of the above disorders should have follow-up enzymatic or molecular testing for confirmation of diagnosis.

Reference Values

CHOLESTANE-3-BETA, 5-ALPHA, 6-BETA-TRIOL

Cutoff: < or =0.070 nmol/mL

7-KETOCHOLESTEROL

Cutoff: < or =0.100 nmol/mL

LYSO-SPHINGOMYELIN

Cutoff: < or =0.100 nmol/mL

GLUCOPSYCHOSINE

Cutoff: < or =0.003 nmol/mL

7-ALPHA-HYDROXY-4-CHOLESTEN-3-ONE (7a-C4)

Cutoff: < or =0.300 nmol/mL

7-ALPHA,12-ALPHA-DIHYDROXYCHOLEST-4-en-3-ONE (7a12aC4)

Cutoff: < or =0.100 nmol/mL

Interpretation

An elevation of 7-alpha-hydroxy-4-cholesten-3-one (7a-C4) and 7-alpha,12-alpha-dihydroxycholest-4-en-3-one (7a12aC4) is strongly suggestive of cerebrotendinous xanthomatosis .

An elevation of glucopsychosine is indicative of Gaucher disease.

An elevation of lyso-sphingomyelin (LSM) and LSM 509 is highly suggestive of Niemann-Pick type A or B disease.

An elevation of cholestane-3-beta, 5-alpha, 6-beta-triol and LSM 509 is highly suggestive of Niemann-Pick disease type C.

Cautions

Patients with Wolman disease or cholestatic biliary atresia may have a profile similar to Niemann-Pick disease type C.

Patients with bile acid malabsorption or ileal resection may have elevations of 7-alpha-hydroxy-4-cholesten-3-one (7aC4).

This test does not identify all causes of hepatosplenomegaly.

A positive test result is strongly suggestive of a diagnosis but needs follow-up by stand-alone biochemical or molecular assay.

Clinical Reference

1. DeBarber AE, Luo J, Star-Weinstock M, et al: A blood test for cerebrotendinous xanthomatosis with potential for disease detection in newborns. *J. Lipid Res.* 2014 Jan;55(1):146-154
2. Federico A, Dotti MT, Gallus GN: Cerebrotendinous xanthomatosis. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. *GeneReviews* [Internet]. University of Washington, Seattle; 2003. Updated April 14, 2016. Accessed December 14, 2022. Available at www.ncbi.nlm.nih.gov/books/NBK1409/
3. Grabowski GA, Petsko GA, Phil D, Kolodny EH: Gaucher disease. In: Valle DL, Antonarakis S, Ballabio A, Beaudet AL, Mitchell GA, eds. *The Online Metabolic and Molecular Bases of Inherited Disease*. McGraw-Hill; 2019. Accessed December 14, 2022. Available at <https://ommbid.mhmedical.com/content.aspx?sectionid=225546056&bookid=2709>
4. Murugeasan V, Chuan WL, Liu J, et al: Glucosylsphingosine is a key biomarker of Gaucher disease. *Am J Hematol.* 2016 Nov;91(11):1082-1089
5. Wasserstein MP, Schuchman EH. Acid sphingomyelinase deficiency. In: Adam MP, Everman DB, Mirzaa GM, et al., eds. *GeneReviews* [Internet]. University of Washington, Seattle; 2006. Updated February 25, 2021. Accessed December 14, 2022. Available at www.ncbi.nlm.nih.gov/books/NBK1370/
6. Wasserstein M, Dionisi-Vici C, Giugliani R, Hwu WL, Lidove O, Lukacs Z, Mengel E, Mistry PK, Schuchman EH,

McGovern M. Recommendations for clinical monitoring of patients with acid sphingomyelinase deficiency (ASMD). Mol Genet Metab. 2019 Feb;126(2):98-105

7. Patterson M: Niemann-Pick disease type C. In: Adam MP, Everman DB, Mirzaa GM, et al, eds. GeneReviews [Internet]. University of Washington, Seattle; 2000. Updated December 10, 2020. Accessed December 14, 2022. Available at www.ncbi.nlm.nih.gov/books/NBK1296/

8. Geberhiwot T, Moro A, Dardis A, et al; International Niemann-Pick Disease Registry (INPDR): Consensus clinical management guidelines for Niemann-Pick disease type C. Orphanet J Rare Dis. 2018 Apr 6;13(1):50

Performance

Method Description

An internal standard is added to an aliquot of plasma, which is then subjected to protein precipitation. Following centrifugation, the supernatant is subjected to liquid chromatography tandem mass spectrometry (LC-MS/MS) analysis. The MS/MS is operated in the multiple reaction monitoring positive mode to follow the precursor to product species transitions for each analyte and internal standard. The ratio of the extracted peak areas to internal standard determined by the LC-MS/MS is used to calculate the concentration of in the sample.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Tuesday, Thursday

Report Available

3 to 7 days

Specimen Retention Time

2 months

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

82542

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
HSMP	Hepatosplenomegaly Panel, P	92743-4

Result ID	Test Result Name	Result LOINC® Value
601542	Interpretation (HSMP)	59462-2
601536	Cholestane-3beta,5alpha,6beta-triol	92755-8
601537	7-Ketocholesterol	92764-0
601538	Lyso-sphingomyelin	92747-5
601539	Glucopsychosine	92750-9
601540	7a-hydroxy-4-cholesten-3-one	92761-6
601541	7a,12a-dihydroxycholest-4-en-3-one	92758-2
601543	Reviewed By	18771-6