

Overview

Useful For
Classification of lymphomas

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm
[For the initial technical component only immunohistochemical \(IHC\) stain performed, the appropriate bill-only test ID will be reflexed and charged \(IHTOI\). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged \(IHTOA\).](#)

Method Name
Immunohistochemistry (IHC)

NY State Available
Yes

Specimen

Specimen Type
TECHONLY

Ordering Guidance
This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions
Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required
Supplies: Immunostain Technical Only Envelope (T693)
Specimen Type: Tissue
Container/Tube: Immunostains Technical Only Envelope
Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

- 1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/>
- 2. Clients ordering stains using a manual requisition form will not have access to digital images.
- 3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/#FAQ>

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides	Reject
--	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

CD25 (cluster of differentiation 25) is the receptor for IL2 and is expressed on activated T cells, B cells, and macrophages. It will stain only scattered cells in normal tonsil. CD25 is expressed in certain types of B-cell lymphoma (hairy cell leukemia) and T-cell lymphoma (adult T-cell lymphoma/leukemia [ATLL]). An anti-CD25 therapy can be used in patients who have lymphomas that express CD25.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is

required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

1. Kageyama Y, Miwa H, Arakawa R, et al. Expression of CD25 fluctuates in the leukemia-initiating cell population of CD25-positive AML. PLoS One. 2018;13(12):e0209295. doi:10.1371/journal.pone.0209295
2. deLeeuw RJ, Kroeger DR, Kost SE, Chang PP, Webb JR, Nelson BH. CD25 identifies a subset of CD4+FoxP3- TIL that are exhausted yet prognostically favorable in human ovarian cancer. Cancer Immunol Res. 2015;3(3):245-253. doi:10.1158/2326-6066.CIR-14-0146
3. Zhang W, Pan Y, Gou P, et al. Effect of xanthohumol on Th1/Th2 balance in a breast cancer mouse model. Oncol Rep. 2018;39(1):280-288. doi:10.3892/or.2017.6094
4. Liu Y, Zhang H, Wang Z, Wu P, Gong W. 5-Hydroxytryptamine1a receptors on tumour cells induce immune evasion in lung adenocarcinoma patients with depression via autophagy/pSTAT3. Eur J Cancer. 2019;114:8-24. doi:10.1016/j.ejca.2019.03.017
5. Magaki S, Hojat SA, Wei B, So A, Yong WH. An Introduction to the Performance of Immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance**Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary
88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
CD25	CD25 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
70696	CD25 IHC, Tech Only	Bill only; no result