

CD2 Immunostain, Technical Component Only

## Overview

#### **Useful For**

Determining T-cell lineage

#### **Reflex Tests**

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

### **Testing Algorithm**

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

#### **Method Name**

Immunohistochemistry (IHC)

#### **NY State Available**

Yes

## Specimen

## **Specimen Type**

**TECHONLY** 

# **Ordering Guidance**

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

#### **Shipping Instructions**

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

## Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue

**Container/Tube:** Immunostains Technical Only Envelope

Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick.

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block



CD2 Immunostain, Technical Component Only

## **Digital Image Access**

- 1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <a href="https://news.mayocliniclabs.com/ihc-stains/">https://news.mayocliniclabs.com/ihc-stains/</a>
- 2. Clients ordering stains using a manual requisition form will not have access to digital images.
- 3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <a href="https://news.mayocliniclabs.com/ihc-stains/#FAQ">https://news.mayocliniclabs.com/ihc-stains/#FAQ</a>

#### **Forms**

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u>
Request (T763) with the specimen.

## Reject Due To

Wet/frozen	Reject
tissue	
Cytology	
smears	
Nonformalin	
fixed tissue	
Nonparaffin	
embedded	
tissue	
Noncharged	
slides	
ProbeOn slides	

## **Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

# Clinical & Interpretive

#### **Clinical Information**

CD2 is a pan T-cell antigen that is expressed on normal and neoplastic T cells and natural killer cells. CD2 immunostaining is useful in determining T-cell lineage in cases of T-cell lymphoma.

#### Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.



CD2 Immunostain, Technical Component Only

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

#### **Cautions**

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen-dependent. Best practice is for paraffin sections to be cut within 6 weeks.

#### **Clinical Reference**

- 1. Campbell AM, Peters SB, Zirwas MJ, et al. Immunophenotypic diagnosis of primary cutaneous lymphomas. A review for the practicing dermatologist. J Clin Aesthet Dermatol. 2010;3(10):21-25
- 2. Rai MP, Bedi PS, Marinas EB, Khan NNS. Angioimmunoblastic T-cell lymphoma: a rare subtype of peripheral T-cell lymphoma. Clin Case Rep. 2018;6(4):750-752. doi:10.1002/ccr3.1388
- 3. Stenman L, Persson M, Enlund F, Clasen-Linde E, Stenman G, Heegaard S. Primary orbital precursor T-cell lymphoblastic lymphoma: Report of a unique case. Mol Clin Oncol. 2016;5(5):593-595. doi:10.3892/mco.2016.1008
- 4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5\_25

# **Performance**

## **Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

### **PDF Report**

No

#### Day(s) Performed

Monday through Friday

# Report Available

1 to 3 days

#### **Specimen Retention Time**

Until staining is complete

# **Performing Laboratory Location**

Rochester

# **Fees & Codes**

### Fees

• Authorized users can sign in to <u>Test Prices</u> for detailed fee information.



CD2 Immunostain, Technical Component Only

- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

# **Test Classification**

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

# **CPT Code Information**

88342-TC, primary 88341-TC, if additional IHC

### **LOINC®** Information

Test ID	Test Order Name	Order LOINC® Value
CD2	CD2 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
70697	CD2 IHC, Tech Only	Bill only; no result