

CD34 Immunostain, Technical Component
Only

Overview

Useful For

A marker of immaturity in the setting of acute myeloid leukemia or B-cell lymphoblastic leukemia

Diagnosis of gastrointestinal stromal tumors, solitary fibrous tumors, and angiosarcomas

Reflex Tests

| Test Id | Reporting Name | Available Separately | Always Performed |
|---------|---------------------------|----------------------|------------------|
| IHTOI | IHC Initial, Tech Only | No | No |
| IHTOA | IHC Additional, Tech Only | No | No |

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)

Specimen Type: Tissue



CD34 Immunostain, Technical Component
Only

Container/Tube: Immunostain Technical Only Envelope

Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick.

Acceptable: Formalin-fixed, paraffin-embedded (FFPE) tissue block

Digital Image Access

- 1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/
- 2. Clients ordering stains using a manual requisition form will not have access to digital images.
- 3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: https://news.mayocliniclabs.com/ihc-stains/#FAQ

Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u>
Request (T763) with the specimen.

Reject Due To

| Wet/frozen | Reject |
|----------------|--------|
| tissue | |
| Cytology | |
| smears | |
| Nonformalin | |
| fixed tissue | |
| Nonparaffin | |
| embedded | |
| tissue | |
| Noncharged | |
| slides | |
| ProbeOn slides | |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|---------------------|------|-------------------|
| TECHONLY | Ambient (preferred) | | |
| | Refrigerated | | |

Clinical & Interpretive

Clinical Information

CD34 (cluster of differentiation 34) is 115 kDa membrane-associated antigen found on human hematopoietic progenitor cells and vascular endothelial cells. In normal tonsil sections, antibodies to CD34 strongly stain vascular endothelial cells.

CD34 is used as a marker of immaturity in the setting of acute myeloid leukemia or B-cell lymphoblastic leukemia. It is



CD34 Immunostain, Technical Component
Only

also useful in the diagnosis of gastrointestinal stromal tumors, solitary fibrous tumors, and angiosarcomas.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

- 1. Lopez-Nunez O, Surrey LF, Alaggio R, Fritchie KJ, John I. Novel PPP1CB-ALK fusion in spindle cell tumor defined by S100 and CD34 coexpression and distinctive stromal and perivascular hyalinization. Genes Chromosomes Cancer. 2020;59(8):495-499. doi:10.1002/gcc.22844
- 2. Matoso A, Epstein JI. Epithelioid angiosarcoma of the bladder: A series of 9 cases. Am J Surg Pathol. 2015;39(10):1377-1382. doi:10.1097/PAS.0000000000000444
- 3. Stanek J, Abdaljaleel M. CD34 immunostain increases the sensitivity of placental diagnosis of fetal vascular malperfusion in stillbirth. Placenta. 2019;77:30-38. doi:10.1016/j.placenta.2019.02.001
- 4. Sardina LA, Piliang M, Bergfeld WF. Diagnostic value of CD34 and calretinin immunostaining in the diagnosis of proliferating tricholemmal tumor and trichoblastoma. Int J Dermatol. 2020;59(1):99-102. doi:10.1111/ijd.14461
- 5. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days



CD34 Immunostain, Technical Component
Only

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary 88341-TC, if additional IHC

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|---------------------|----------------------|
| CD34I | CD34 IHC, Tech Only | Order only;no result |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|---------------------|----------------------|
| 70701 | CD34 IHC, Tech Only | Bill only; no result |