

Helicobacter pylori (H pylori) Immunostain, Technical Component Only

Overview

Useful For

Aiding in the identification of Helicobacter pylori infection

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain (no pathologist interpretation is performed). If diagnostic consultation by a pathologist is required order PATHC / Pathology Consultation.

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Supplies: Immunostain Technical Only Envelope (T693)
Specimen Type: Tissue
Container/Tube: Immunostain Technical Only Envelope
Preferred: 2 Unstained positively charged glass slide (25- x 75- x 1-mm) per test ordered; sections 4-microns thick



Helicobacter pylori (H pylori) Immunostain, Technical Component Only

Acceptable: Formalin-fixed, paraffin-embedded tissue block

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <u>https://news.mayocliniclabs.com/pathology/digital-imaging/</u>

2. Clients ordering stains using a manual requisition form will not have access to digital images.

3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <u>https://news.mayocliniclabs.com/pathology/digital-imaging/#section3</u>

Forms

If not ordering electronically, complete, print, and send a <u>Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains</u> <u>Request</u> (T763) with the specimen.

Reject Due To

Wet/frozen	Reject
tissue Cytology	
smears	
Nonformalin	
fixed tissue	
Nonparaffin	
embedded	
tissue	
Noncharged	
slides ProbeOn	
slides	

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Helicobacter pylori is a bacterium that frequently infects the stomach, colonizing the gastric pits. *H pylori* infection is associated with the development of gastroduodenal ulcers and gastric mucosa-associated lymphoid tissue (MALT) lymphomas.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.



Helicobacter pylori (H pylori) Immunostain, Technical Component Only

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

Clinical Reference

 Patnayak R, Reddy V, Jena A, et al. Helicobacter pylori in cholecystectomy specimens-morphological and immunohistochemical assessment. J Clin Diagn Res. 2016;10(5):EC01-3. doi:10.7860/JCDR/2016/14802.7716
Son JH, Lebwohl B, Sepulveda AR, Lagana SM. Utilization rate of Helicobacter pylori immunohistochemistry is not associated with the diagnostic rate of helicobacter pylori infection. Appl Immunohistochem Mol Morphol. 2019;27(9):694-698. doi:10.1097/PAI.00000000000680

3. Lee JY, Kim N. Diagnosis of Helicobacter pylori by invasive test: histology. Ann Transl Med. 2015;3(1):10. doi:10.3978/j.issn.2305-5839.2014.11.03

4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed Monday through Friday

Report Available 1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Rochester



Helicobacter pylori (H pylori) Immunostain, Technical Component Only

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test was developed using an analyte specific reagent. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

88342-TC, primary 88341-TC, if additional IHC

LOINC[®] Information

Test ID	Test Order Name	Order LOINC [®] Value
HPYL	H. pylori IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC [®] Value
70779	H. pylori IHC, Tech Only	Bill only; no result