# Test Definition: TCL1A

T-Cell Leukemia/Lymphoma Protein 1A (TCL1A) Immunostain, Technical Component Only

## Overview

### Useful For

Identification of T-cell leukemia and lymphoma protein overexpression in neoplasms

### Reflex Tests

| Test Id | Reporting Name | Available Separately | Always Performed |
|---|---|---|---|
| IHTOI | IHC Initial, Tech Only | No | No |
| IHTOA | IHC Additional, Tech Only | No | No |

### Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

### Method Name

Immunohistochemistry (IHC)

### NY State Available

Yes

## Specimen

### Specimen Type

TECHONLY

### Ordering Guidance

This test includes only technical performance of the stain; **no pathologist interpretation is provided**.

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled**. Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

### Shipping Instructions

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

### Specimen Required

# Test Definition: TCL1A

T-Cell Leukemia/Lymphoma Protein 1A (TCL1A) Immunostain, Technical Component Only

**Specimen Type:** Tissue
**Supplies:** Immunostain Technical Only Envelope (T693)
**Container/Tube:** Immunostain Technical Only Envelope
**Submit:**
-Formalin-fixed, paraffin-embedded tissue block
OR
-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick
**Additional Information**: Broken or cracked unstained slides will be rejected. Stained slides that were broken and repaired (ie, double mounted) at the client site will not be rejected, as, in most cases, they can still be reviewed under the microscope. However, due to the double mounting, digital scanning cannot be performed.

## Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/pathology/digital-imaging/
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: https://news.mayocliniclabs.com/pathology/digital-imaging/#section3

## Forms

If not ordering electronically, complete, print, and send a Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains Request (T763) with the specimen.

## Reject Due To

| | |
|---|---|
| Wet/frozen tissue | Reject |
| Cytology smears | Reject |
| Nonformalin fixed tissue | Reject |
| Nonparaffin embedded tissue | Reject |
| Noncharged slides | Reject |
| ProbeOn slides | Reject |
| Snowcoat slides | Reject |
| Broken or cracked unstained slides | Reject |
| Broken or | Reject |

| | |
|---|---|
| cracked stained slides that cannot be repaired | |

### Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---|---|---|---|
| TECHONLY | Ambient (preferred) | | |
| | Refrigerated | | |

## Clinical & Interpretive

### Clinical Information
T-cell leukemia/lymphoma (TCL) proteins augment AKT signal transduction and enhance cell proliferation and survival. Inversion or translocations involving the *TCL-1* gene are present in more than 90% of T-cell prolymphocytic leukemias, resulting in overexpression of the TCL-1 protein. TCL-1 is also found in plasmacytoid monocytes in reactive tissues, and the putative malignant counterpart, blastic plasmacytoid dendritic cell neoplasm. In normal tonsil, expression is limited to the B-cell compartment.

### Interpretation
This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

### Cautions
Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:
-Minimize time slides are stored after being unpackaged
-Limit exposure to high humidity and heat
-Minimize exposure to plastics

Broken or cracked unstained slides are not compatible with the instrumentation used in our performing laboratories and

will be rejected. Additional material may be requested.

Attempts will be made to repair broken or cracked stained slides; however, if they are unable to be repaired (ie, shattered in transit), they will be rejected and discarded due to safety concerns.

### Clinical Reference

1.Said JW, Hoyer KK, French SW, et al. TCL1 oncogene expression in B cell subsets from lymphoid hyperplasia and distinct classes of B cell lymphoma. Lab Invest. 2001;81(4):555-564
2. Herling M, Teitell MA, Shen RR, et al. TCL1 expression in plasmacytoid dendritic cells (DC2s) and the related CD4+ CD56+ blastic tumors of skin. Blood. 2003;101(12):5007-5009
3. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

## Performance

### Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

### PDF Report

No

### Day(s) Performed

Monday through Friday

### Report Available

1 to 3 days

### Specimen Retention Time

Until staining is complete

### Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

# Test Definition: TCL1A

T-Cell Leukemia/Lymphoma Protein 1A (TCL1A) Immunostain, Technical Component Only

## Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

## LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---|---|---|
| TCL1A | TCL1A IHC, Tech Only | Order only;no result |

| Result ID | Test Result Name | Result LOINC® Value |
|---|---|---|
| 70873 | TCL1A IHC, Tech Only | Bill only; no result |