MAYO CLINIC LABORATORIES

Test Definition: TRAP

Tartrate-Resistant Acid Phosphatase (TRAP)
Immunostain, Technical Component Only

Overview

Useful For
Classification of leukemias and lymphomas

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm
For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name
Immunohistochemistry (IHC)

NY State Available
Yes

Specimen

Specimen Type
TECHONLY

Ordering Guidance
This test includes only technical performance of the stain; **no pathologist interpretation is provided**.

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled**. Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

Shipping Instructions
Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required
Specimen Type: Tissue

Test Definition: TRAP

Tartrate-Resistant Acid Phosphatase (TRAP)
Immunostain, Technical Component Only

Supplies: Immunostain Technical Only Envelope (T693)
Container/Tube: Immunostain Technical Only Envelope
Submit:
-Formalin-fixed, paraffin-embedded tissue block
OR
-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick
Additional Information: Broken or cracked unstained slides will be rejected. Stained slides that were broken and repaired (ie, double mounted) at the client site will not be rejected, as, in most cases, they can still be reviewed under the microscope. However, due to the double mounting, digital scanning cannot be performed.

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: https://news.mayocliniclabs.com/pathology/digital-imaging/
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: https://news.mayocliniclabs.com/pathology/digital-imaging/#section3

Forms

If not ordering electronically, complete, print, and send a Immunohistochemical (IHC)/In Situ Hybridization (ISH) Stains Request (T763) with the specimen.

Reject Due To

Wet/frozen tissue	Reject
Cytology smears	Reject
Nonformalin fixed tissue	Reject
Nonparaffin embedded tissue	Reject
Noncharged slides	Reject
ProbeOn slides	Reject
Snowcoat slides	Reject
Broken or cracked unstained slides	Reject
Broken or cracked stained slides	Reject

that cannot be repaired	

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

Tartrate-resistant acid phosphatase (TRAP) is a basic, iron-binding protein found within the cytoplasmic granules of cells of hairy cell leukemia. TRAP can be useful in distinguishing hairy cell leukemia from other types of B-cell lymphomas. Caution should be used since other types of lymphoma can also be positive, such as marginal zone lymphoma, but usually their staining is less intense. Normal mast cells also express TRAP.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:
-Minimize time slides are stored after being unpackaged
-Limit exposure to high humidity and heat
-Minimize exposure to plastics

Broken or cracked unstained slides are not compatible with the instrumentation used in our performing laboratories and will be rejected. Additional material may be requested.

Attempts will be made to repair broken or cracked stained slides; however, if they are unable to be repaired (ie, shattered in transit), they will be rejected and discarded due to safety concerns.

Clinical Reference

1. Dunphy CH. Reaction patterns of TRAP and DBA.44 in hairy cell leukemia, hairy cell variant, and nodal and extranodal marginal zone B-cell lymphomas. Appl Immunohistochem Mol Morphol. 2008;16(2):135-139
2. Hoyer JD, Li CY, Yam LT, Hanson CA, Kurtin PJ. Immunohistochemical demonstration of acid phosphatase isoenzyme 5 (tartrate-resistant) in paraffin sections of hairy cell leukemia and other hematologic disorders. Am J Clin Pathol. 1997;108(3):308-315
3. Janckila AJ, Cardwell EM, Yam LT, Li CY. Hairy cell identification by immunohistochemistry of tartrate-resistant acid phosphatase. Blood. 1995;85(10):2839-2844
4. Janckila AJ, Slone SP, Lear SC, Martin A, Yam LT. Tartrate-resistant acid phosphatase as an immunohistochemical marker for inflammatory macrophages. Am J Clin Pathol. 2007;127(4):556-566
5. Sherman MJ, Hanson CA, Hoyer JD. An assessment of the usefulness of immunohistochemical stains in the diagnosis of hairy cell leukemia. Am J Clin Pathol. 2011;136(3):390-399
6. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. Methods Mol Biol. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.

- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary
88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
TRAP	TRAP IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
70882	TRAP IHC, Tech Only	Bill only; no result