MAYO CLINIC LABORATORIES

Test Definition: FMARC

Bupivacaine (Marcaine)

Overview

Method Name
Liquid Chromatography/ Tandem Mass Spectrometry (LC/MS/MS)

NY State Available
Yes

Specimen

Specimen Type
Serum Red

Specimen Required
Draw blood in a plain red-top tube(s), **serum gel tube is not acceptable**. Spin down and send 1 mL of serum frozen in a plastic vial.

Specimen Minimum Volume
0.5 mL

Reject Due To

Hemolysis	NA
Lipemia	NA
Icterus	NA
Other	NA

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Frozen (preferred)	180 days	
	Ambient	72 hours	
	Refrigerated	72 hours	

Clinical & Interpretive

Reference Values
Reference Range: 0.1 - 4.0 ug/mL

Test Definition: FMARC

Bupivacaine (Marcaine)

Performance

PDF Report
No

Day(s) Performed
Monday through Sunday

Report Available
5 to 11 days

Performing Laboratory Location
Medtox Laboratories, Inc.

Fees & Codes

Fees
- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification
This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

CPT Code Information
80299

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FMARC	Bupivacaine (Marcaine)	3412-4

Result ID	Test Result Name	Result LOINC® Value
Z5052	Bupivacaine	3412-4