MAYO CLINIC LABORATORIES

Test Definition: FPDD

Pigeon/Dove Droppings Gel Diffusion

Overview

Method Name
Gel Immunodiffusion

NY State Available
Yes

Specimen

Specimen Type
Serum

Specimen Required
Specimen Type: Serum
Container/Tube: Red or SST
Specimen Volume: 0.5 mL

Collection Instructions: Draw blood in a plain red-top tube(s), serum gel tube(s) is acceptable. Spin down and send 0.5 mL of serum refrigerated in a plastic vial.

Specimen Minimum Volume
0.5 mL

Reject Due To

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	28 days	
	Ambient	28 days	
	Frozen	28 days	

Clinical & Interpretive

Reference Values
Negative

Interpretation

MAYO CLINIC LABORATORIES

Test Definition: FPDD

Pigeon/Dove Droppings Gel Diffusion

The gel diffusion method was used to test this patient's serum for the presence of precipitating antibodies (IgG) to the antigens indicated. These antibodies are serological markers for exposure and immunological sensitization. The clinical significance varies, depending on the history and symptoms.

Performance

PDF Report
No

Day(s) Performed
Tuesday, Friday, Saturday

Report Available
7 to 14 days

Performing Laboratory Location
Eurofins Viracor

Fees & Codes

Fees
- Authorized users can sign in to Test Prices for detailed fee information.
- Clients without access to Test Prices can contact Customer Service 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact Customer Service.

Test Classification
This test was developed and its performance characteristics determined by Eurofins Viracor. It has not been cleared or approved by the U.S. Food and Drug Administration.

CPT Code Information
86331

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
FPDD	Pigeon/Dove Droppings Gel Diffusion	24505-0

Result ID	Test Result Name	Result LOINC® Value
Z5608	Pigeon/Dove Droppings Gel Diffusion	24505-0