

Test Definition: FHV8P

Herpesvirus 8 (HHV-8) DNA, Quantitative Real-Time PCR

Overview

Method Name

Real-Time Polymerase Chain Reaction, RT-PCR

NY State Available

Yes

Specimen

Specimen Type

Varies

Specimen Required

Submit only 1 of the following specimens:

Whole Blood

Specimen Type: Whole Blood

Container/Tube: Lavender-top (EDTA) or yellow-top (ACD)

Specimen Volume: 1 mL

Collection Instructions: Draw blood in a lavender-top (EDTA) tube(s) or yellow-top (ACD) tube(s) and send 1 mL whole

blood refrigerated.

Stability: Ambient 48 hours; Refrigerated 7 days: Frozen 30 days

Serum

Specimen Type: Serum

Collection Container/Tube: Red-top

Submission Container/Tube: 12x75 mm screw-capped vial

Specimen Volume: 1 mL

Collection Instructions: Draw blood in a plain red-top tube(s). Spin down and send 1 mL serum in a plastic,

screw-capped vial. Send specimen refrigerated.

Stability: Ambient 48 hours; Refrigerated 7 days, Frozen 30 days

Plasma

Collection Container/Tube: yellow-top (ACD), lavender-top (EDTA), or PPT (white-top) tube

Submission Container/Tube: 12x57 mm screw-capped vial

Specimen Volume: 1 mL



Test Definition: FHV8P

Herpesvirus 8 (HHV-8) DNA, Quantitative Real-Time PCR

Collection Instructions: Draw blood in a yellow-top (ACD) or lavender-top (EDTA) tube(s). Spin down and transfer 1 mL ACD or EDTA plasma into a plastic, screw-capped vial. Send specimen refrigerated.

Stability: Ambient 48 hours; Refrigerated 7 days; Frozen 30 days

Specimen Minimum Volume

0.5 mL

Reject Due To

| Thawing: | Cold OK; Warm reject |
|---------------|---|
| Other reasons | Sodium heparin or Lithium heparin tubes |
| for rejection | |

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|--------------------------|----------|-------------------|
| Varies | Refrigerated (preferred) | 7 days | |
| | Frozen | 30 days | |
| | Ambient | 48 hours | |

Clinical & Interpretive

Clinical Information

Herpesvirus 8 (HHV-8) DNA, Quantitative Real-Time PCR-Human herpesvirus-8 (HHV-8) is associated with the development of all forms of Kaposi's sarcoma, as well as some other rare lymphoproliferative diseases, such as primary effusion lymphoma and multicentric Castleman's disease. Quantitative PCR may be used to monitor the level of viremia in a patient, often in the context of therapy.

Reference Values

Not Detected

Performance

PDF Report

No

Day(s) Performed

Monday through Sunday

Report Available

3 to 6 days



Test Definition: FHV8P

Herpesvirus 8 (HHV-8) DNA, Quantitative Real-Time PCR

Performing Laboratory Location

Quest Diagnostics

Fees & Codes

Fees

- Authorized users can sign in to <u>Test Prices</u> for detailed fee information.
- Clients without access to Test Prices can contact <u>Customer Service</u> 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact <u>Customer Service</u>.

Test Classification

This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics. It has not been cleared or approved by FDA. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes.

CPT Code Information

87799

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|----------------------------------|--------------------|
| FHV8P | Herpes Virus 8 DNA, Quant RT-PCR | 49406-2 |

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|---------------------------|---------------------|
| Z6082 | Source | 31208-2 |
| Z6083 | Herpesvirus 8 DNA, QN PCR | 49406-2 |
| Z6084 | Herpesvirus 8 DNA, QN PCR | No LOINC Needed |