



# Test Definition: FBCFS

Barbiturates Confirmation, Serum

## Overview

### Method Name

Only orderable as a reflex. For more information see:  
FDA1S / Drugs of Abuse (10 panel) and Alcohol Screen, Serum  
FD10S / Drugs of Abuse Screen, Serum

Gas Chromatography/Mass Spectrometry (GC/MS)

### NY State Available

Yes

## Specimen

### Specimen Type

Serum Red

### Specimen Required

Only orderable as a reflex. For more information see:  
FDA1S / Drugs of Abuse (10 panel) and Alcohol Screen, Serum  
FD10S / Drugs of Abuse Screen, Serum

### Reject Due To

All specimens will be evaluated by the processing and performing laboratories for test suitability

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum Red	Refrigerated (preferred)	14 days	
	Frozen	30 days	

## Clinical & Interpretive

### Clinical Information

[Refer to www.nmslabs.com/test-catalog](http://www.nmslabs.com/test-catalog)

### Reference Values

Only orderable as a reflex. For more information see:  
FDA1S / Drugs of Abuse (10 panel) and Alcohol Screen, Serum  
FD10S / Drugs of Abuse Screen, Serum

Reporting limit determined each analysis.

Butabarbital: None detected  
Butalbital: None detected  
Pentobarbital: None detected  
Secobarbital: None detected  
Phenobarbital: None detected

**Interpretation**

Butabarbital:

Plasma concentrations of 2-3 mcg/mL produce sedation and plasma concentrations of 25 mcg/mL produce sleep in most patients. Plasma concentrations of greater than 30 mcg/mL may produce coma and plasma concentrations in excess of 50 mcg/mL are potentially lethal.

Butalbital:

A single oral 100 mcg dose resulted in a mean peak blood concentration of 2.1 mcg/mL (range: 1.7-2.6 mcg/mL) at 2 hours, with a decline to 1.5 mcg/mL (range: 1.3-1.7 mcg/mL) by 24 hours. Potentially toxic at plasma concentrations greater than 10 mcg/mL.

Pentobarbital:

Peak serum concentrations of 1.2-3.1 mcg/mL were produced 0.5-2.0 hours after a 100 mg oral dose and peak serum concentration of 3 mcg/mL were produced 6 min. following a 100 mg IV dose. Potentially toxic at blood concentrations greater than 10 mcg/mL.

Secobarbital:

A 3.3 mg/kg oral dose (approximately 230 mg/70 kg) produced a mean peak blood concentration of 2.0 mcg/mL (range, 1.8-2.2 mcg/mL) at 3 hours, diminishing to 1.3 mcg/mL by 20 hours and 0.8 mcg/mL by 40 hours. Potentially toxic at blood concentrations greater than 8 mcg/mL.

Phenobarbital:

Recommended serum concentration range during anticonvulsant therapy with primidone: 10-40 mcg/mL.

**Performance****PDF Report**

No

**Performing Laboratory Location**

NMS Labs

**Fees & Codes**

**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**CPT Code Information**

80345

G0480 (if appropriate)

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
FBCFS	Barbiturates Confirmation, S	Not Provided

Result ID	Test Result Name	Result LOINC® Value
Z5287	Butabarbital	18384-8
Z5288	Butalbital	82971-3
Z5290	Pentobarbital	82969-7
Z5291	Secobarbital	82968-9
Z5292	Phenobarbital	60468-6