

Overview

Useful For

Aiding in the diagnosis of Eastern equine encephalitis using serum specimens

Testing Algorithm

For more information see [Mosquito-borne Disease Laboratory Testing](#).

Special Instructions

- [Mosquito-borne Disease Laboratory Testing](#)

Method Name

Immunofluorescence Assay (IFA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions: Centrifuge and aliquot serum into plastic vial.

Forms

If not ordering electronically, complete, print, and send [Infectious Disease Serology Test Request](#) (T916) with the specimen.

Specimen Minimum Volume

0.15 mL

Reject Due To

Gross	Reject
-------	--------

hemolysis	
Gross lipemia	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	14 days	
	Frozen	14 days	

Clinical & Interpretive

Clinical Information

Eastern equine encephalitis (EEE) is within the alphavirus group. It is a low prevalence cause of human disease in the Eastern and Gulf Coast states. EEE is maintained by a cycle of mosquito/wild bird transmission, peaking in the summer and early fall, when humans may become an adventitious host. The most common clinically apparent manifestation is a mild undifferentiated febrile illness, usually with headache.

Central nervous system involvement is demonstrated in only a minority of infected individuals; it is more abrupt and more severe with EEE than other arboviruses, with children being more susceptible to severe disease. Fatality rates are approximately 70% for EEE.

Reference Values

IgG: <1:10
IgM: <1:10
Reference values apply to all ages.

Interpretation

In patients infected with this virus, IgG antibody is generally detectable within 1 to 3 weeks of onset, peaking within 1 to 2 months, and declining slowly thereafter.

IgM class antibody is also reliably detected within 1 to 3 weeks of onset, peaking and rapidly declining within 3 months.

Single serum specimen IgG greater than or equal to 1:10 indicates exposure to the virus.

Results from a single serum specimen can differentiate early (acute) infection from past infection with immunity if IgM is positive (suggests acute infection).

A 4-fold or greater rise in IgG antibody titer in acute and convalescent sera indicate recent infection.

In the United States it is unusual for any patient to show positive reactions to more than 1 of the arboviral antigens, although Western equine encephalitis and Eastern equine encephalitis antigens will show a noticeable cross-reactivity.

Infections can occur at any age. The age distribution depends on the degree of exposure to the particular transmitting

arthropod relating to age and sex, as well as the occupational, vocational, and recreational habits of the individuals. Once humans have been infected, the severity of the host response may be influenced by age.

Cautions

All results must be correlated with clinical history and other data available to the attending physician.

Specimens collected within the first 2 weeks after onset are variably negative for IgG antibody and should not be used to exclude the diagnosis of arboviral disease. If arboviral infection is suspected, a second specimen should be collected and tested 10 to 21 days later.

Eastern equine encephalitis and Western equine encephalitis viruses show some cross-reactivity; however, antibody response to the infecting virus is typically at least 8-fold higher.

Usually, when an infection with an arbovirus is suspected, it is too late to isolate the virus or draw serum specimens to detect a rise of antibody titer.

Clinical Reference

1. Gonzalez-Scarano F, Nathanson N: Bunyaviruses. In: Fields BN, Knipe DM, eds. Fields Virology. Vol 1. 2nd ed. Raven Press; 1990:1195-1228
2. Donat JF, Rhodes KH, Groover RV, Smith TF: Etiology and outcome in 42 children with acute nonbacterial meningoencephalitis. Mayo Clin Proc. 1980 Mar;55(3):156-160
3. Tsai TF: Arboviruses. In: Murray PR, Baron EJ, Pfaller MA, et al, eds. Manual of Clinical Microbiology. 7th ed. American Society for Microbiology; 1999:1107-1124
4. Calisher CH: Medically important arboviruses of the United States and Canada. Clin Microbiol Rev. 1994 Jan;7(1):89-116
5. Markoff L: Alphaviruses (Chikungunya, Eastern equine encephalitis). In: Bennett JE, Dolin R, Blaser MJ, eds. Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases. 9th ed. Elsevier; 2020:1997-2006

Performance**Method Description**

The indirect immunofluorescent antibody (IFA) assay is a 2-stage “sandwich” procedure. In the first stage, the patient serum is diluted in Pretreatment Diluent for IgM and phosphate buffered saline (PBS) for IgG, added to appropriate slide wells in contact with the substrate, and incubated. Following incubation, the slide is washed in PBS, which removes unbound serum antibodies. In the second stage, each antigen well is overlaid with fluorescein-labeled antibody to IgM and IgG. The slide is incubated allowing antigen-antibody complexes to react with the fluorescein-labeled anti-IgM and anti-IgG. After the slide is washed, dried, and mounted, it is examined using fluorescence microscopy. Positive reactions appear as cells exhibiting bright apple-green cytoplasmic fluorescence against a background of red negative control cells. Semi-quantitative endpoint titers are obtained by testing serial dilutions of positive specimens.(Package inserts: Arbovirus IFA IgM and Arbovirus IFA IgG Instructions for Use. Focus Diagnostics; Rev. 02, 05/01/2018)

PDF Report

No

Day(s) Performed

May through October: Monday through Friday
November through April: Monday, Wednesday, Friday

Report Available

Same day/1 to 4 days

Specimen Retention Time

2 weeks

Performing Laboratory Location

Rochester

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

86652 x 2

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
EEEE	East Equine Enceph Ab, IgG and IgM, S	69034-7

Result ID	Test Result Name	Result LOINC® Value
83354	East Equine Enceph Ab, IgG, S	In Process
83355	East Equine Enceph Ab, IgM, S	In Process